

Household and Structural Pest Management for Professionals

See *The UT E&PP Redbook, Household and Structural Pests section* (<https://ag.tennessee.edu/EPP/Pages/Household-Structural.aspx>) for information related to preventing and otherwise managing pests using integrated pest management. More information for the pest management professional can be found in the following UT Extension Publications <https://ag.tennessee.edu/PSEP/Pages/studymanuals.aspx>

- Vail, K.M., G. Burgess, R. Gerhardt and C. Harper [eds.]. 2007. PB1673 General Pest and Rodent Control Pesticide Applicator Licensing Manual (GRC). pp. 130.
- Vail, K., G. Burgess, R. Gerhardt and C. Jones [eds.]. 2002. PB1685 Public Health Mosquito Control: The Tennessee Mosquito Control Handbook (Pesticide Applicator Licensing Manual). pp. 54.
- Vail, K., G. Burgess, C. Pless and B. Bonds [eds.]. 2006. PB 1703 Wood-Destroying Organisms Licensing Manual.
- Vail, K.M., E. Burgess, R. Gerhardt, and Craig Harper. 2006. PB1732 Industrial, Institutional, Structural and Health Related Pest Management Certification Manual (Category 7). pp. 105
- Vail, K.M., E.E. Burgess, R. Gerhardt, C. Jones, J. Skinner and C. Harper. 2003. PB1733. Public Health Pest Control (Certification Category 8). pp. 99

Further references for household pest identification:

Manuals:

- Bennett, G. W., J. M. Owens, & R. M. Corrigan. 2011. Scientific Guide to Pest Management Operations, seventh edition Cleveland, OH: Questex Publishing.
- Mallis, A. 2011. **Handbook of Pest Control - the Behavior, Life History and Control of Household Pests.** 10th edition. Mallis Handbook LLC.

Field Guides

- Hedges, S. 2010. **Pest Control Technology Field Guide for the Management of Structure-Infesting Ants, 3rd edition.** G.I.E. Inc. Publishers, Richfield, Ohio.
- Hedges, S. **Pest Control Technology Field Guide for the Management of Structure-Infesting Flies.** Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. 2012. **Pest Control Technology Field Guide for the Management of Urban Spiders.** Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. and M. Lacey. **Structure-Infesting Beetles. Volume 1: Hide and Carpet Beetles/Wood-Boring Beetles.** Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. and M. Lacey. **Structure-Infesting Beetles. Volume 2: Stored Product Beetles/Occasional and Overwintering Beetles.** Franzak and Foster Co.
- Klotz, J., L. Hansen, R. Pospischil and M. Rust. 2008. **Urban Ants of North America and Europe: Identification, Biology and Management.** Comstock Publishing Associates, Cornell University Press, Ithaca, New York pp. 196
- Smith, E and R. Whitman. 2007. **NPMA Field Guide to Structural Pests.** Second edition. <http://nmpapestworld.org/>

General identification guides for insects.

- Petersons Field Guides: Insects of North America # 19, Beetles of North America #29, Moths of Eastern North America #30

The following pages list recommended procedures and, if necessary, pesticides to manage specific pests. Remember to reduce the pest's access to food, water and shelter. If pesticides are needed, they are more likely to provide control if access to these necessities are limited. Additional publications (SPs, PBs) are listed on our web site at <http://eppserver.ag.utk.edu/personnel/Vail/publications.html>. Percentages that appear after the Trade Name indicate percentage of the active ingredient. Where no percentage is given, see the label for more details. SEE THE LABEL FOR ALL DIRECTIONS!

Extensive changes to pyrethroid labels occurred recently to help limit pyrethroid contamination of water and habitat of protected species. An extensive summary of the changes can be found at <http://library.constantcontact.com/download/get/file/1102861932245-488/Letter+to+Registrants+-+January+2013%5B1%5D.pdf>


EPA has recently revised outdoor foliar use directions for products containing the active ingredients imidacloprid, dinotefuran, clothianidin or thiamethoxam with pollinator protection language (<http://www2.epa.gov/pollinator-protection>) and ASPCRO has clarified the language as it pertains to non-agricultural use (<http://pesticidestewardship.org/PollinatorProtection/Documents/ASPCRO%20non-ag%20neonic%20guidance%208-18-14%20final.pdf>). To help pest management professionals effectively control pests while protecting pollinators, NPMA has developed *Pollinator Awareness Training* (pestworld.learningzen.com), pollinator information to share with clientele www.pollinatorhealth.org, and a draft of best management practices for these situations. Additional label language changes to protect pollinators are expected.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS - ARGENTINE ANT, ODOROUS HOUSE ANT</p> <p>PB1629</p>	<p>ARGENTINE ANT Bump (Node) on waist: one, obvious Gaster tip: no circular opening Odor when crushed: "disagreeable, rotten-coconut-like" with an additional faint musty odor Color: light to dark brown</p> <p>ODOROUS HOUSE ANT Nodes on waist: one, very flat, barely noticeable node hidden by the gaster Gaster tip: no circular opening Odor when crushed: "disagreeable, rotten-coconut-like" Color: brown to black Both ant workers: 1/8 inch</p> <p>The odorous house ant is the most common house-invading ant in TN. It nests in shady, moist areas such as under mulch, pine straw, stones, and logs. Moves indoors during periods of heavy rain and moves nests often. Winged male odorous house ants may be collected from lights from May through June.</p> <p>Argentine ant is not as common as the odorous house ant, but can build up larger populations. Behavior is similar for both species.</p>	<p><u>Perimeter Spray:</u> Termidor SC 0.06% Arilon 0.05%, 0.10% Optigard Flex 0.1% (to perimeter and plants combined with Optigard Ant Gel Bait in outdoor stations</p> <p><u>Outdoor Baiting:</u> Advion Insect Granule (perimeter broadcast) Intice Thiquid 1% Gourmet Ant Bait Liquid Maxforce Quantum Ant Bait MAXFORCE@COMPLETE Brand Granular insect Bait Maxforce Ant Killer Bait Gel (for Argentine ant) Niban FG, G baits Optigard Ant Gel Bait PT 388B Advance Ant Gel Bait Terro-PCO (1%)</p> <p><u>Indoor Crack and Crevice Spray and Outdoor Entry Points</u> Phantom</p>	<p>Our best research results for odorous house ants have been a 0.06% fipronil (Termidor SC) spray to entry points and 1 foot up and 1 foot out from the foundation base COMBINED with (1) a liquid or other bait placed in the landscape near/around the structure where ants are active OR COMBINED with (2) Phantom applied indoors as crack and crevice in areas of activity as well as at potential entry points. Gel baits may work longer outdoors when placed in stations. Baits listed under odorous house ant and outdoor baiting have eliminated small, laboratory-maintained odorous house ant colonies within 8 weeks of bait placement or have proven effective in field studies.</p> <p>Ants nesting in structures through the winter can be quite challenging to control and to the pest management professionals' reputation. Often the ant centers of activity are difficult to define because small numbers of ants are found in many different locations. You bait one area and activity ceases, only to be found in another area. It's worth exploring the use of placing Quantum or any other liquid/gel bait in indoor stations near ant activity where residents will not see the station or trailing ants. Moisture is fairly limiting this time of year when indoor heat is running longer because of low outdoor temperatures and liquid/gel bait in a station may serve as moisture source.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS - CARPENTER PB1599</p>	<p>Large, black, red, or red-and-black ants that nest in damp wood. Wingless workers 1/4 to 3/8 inch long. Winged male and female reproductives will fly from a colony. Carpenter ants do not eat wood, but excavate smooth galleries in the wood to raise their young. Piles of coarse sawdust or splintered wood often mixed with insect parts may indicate a carpenter ant nest nearby.</p>	<p><u>Outdoor perimeter spray</u> Termidor SC Arilon Transport GHP</p> <p><u>Baits:</u> <u>Indoors in cracks and crevices where ants are seen:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Maxforce Carpenter Ant Bait Niban FG, G</p> <p><u>Outdoors:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Gourmet Ant Bait Liquid Maxforce Carpenter Ant Bait Gel Maxforce Granular Bait Niban FG, G Optigard Ant Gel Bait Terro-PCO (1%)</p> <p><u>Dusts:</u> Deltadust Boric acid dusts Tempo 1D</p> <p><u>Sprays (or foam according to label):</u> TalstarP, Tempo SC Ultra, Tempo Ultra WP, Cy-Kick, Transport, PT221L, Suspend SC, Bora-Care, Shell-guard, Tim-bor (dust, spray or foam), Armor-Guard (dust, spray or foam)</p>	<ol style="list-style-type: none"> 1. Correct moisture problem, repair leaks and ventilate. 2. If ants entering from or foraging outdoors, spray a slower acting, nonrepellent insecticide around the perimeter with an emphasis on areas where ants are entering the structure. 3. Place baits where ants are actively foraging. If baiting, do not spray ants, trails or baits with a fast-acting, repellent spray because it will kill ants too quickly and will stop them from feeding on the baits. 4. Find and treat (dust, spray or foam) nests in wood parts. <p>Drilling into the wood may be necessary. Dust nests in wall voids. Do not apply sprays near electricity.</p>
<p>ANTS-PHARAOH also called "sugar ants" or "piss ants" PB 1629</p>	<p>Nodes on waist: two Gaster tip: sting Worker size: 1/16 inch Antennae: 12-segmented antennae with a three-segmented club Color: yellow or orange with the end of the abdomen darkened</p> <p>Nests rarely found outdoors; however, almost any indoor crack and crevice close to sources of warmth and water.</p> <p>These ants do not swarm. Colonies multiply by "budding," in which part of an existing colony migrates, carrying brood with or without a queen to a new nesting site. Hundreds of queens and 10,000 - 100,000s workers may be present.</p>	<p><u>Baits:</u> Advion Ant Bait Arenas Maxforce FC Ant Bait Station Niban FG</p> <p><u>Indoor crack and crevice spray and entry points:</u> Phantom SC</p> <p><u>Outdoor entry points and trails around structure</u> Termidor SC</p>	<p>Because Pharaoh ant colonies are hidden and can occur in virtually any crack or crevice, baiting is the best way to get an insecticide back to the colony. Give a taste test of baits. Prebait entire structure with honey. Place a bait wherever ants are found. All queen(s) and all immatures must be killed to eliminate a colony. Spraying fast-acting insecticides for Pharaoh ants indoors often worsens the problem by causing the colony to split into many smaller colonies. Apply slower -acting sprays (Phantom, Termidor) around entry points during warm weather when ants are foraging outdoors.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS-FIRE (see Redbook recommendations for Home Lawn Insects). Also see web sites http://www.extension.org/fire_ants for publications (Managing Fire Ants in Urban Areas, Two-step...) and other information about fire ants and "Imported Fire Ants In Tennessee" at http://fireants.utk.edu for extensive list of TN fire ant management products under "Updates". For a list of "homeowner" products by price and application type (broadcast vs. individual mound treatment) see http://www.aces.edu/pubs/docs/A/ANR-0175-A.pdf.</p>			
<p>ASIAN NEEDLE ANT http://eppserver.ag.utk.edu/Whats/wh2011/issue-14-2011.pdf</p>	<p>Both workers and female winged forms can sting, especially when trapped against clothing. Workers (1/5 inch) and queens (1/4 inch) are dark brown, with the legs, mandibles, and outer antennal segments slightly lighter. Workers with a middle, side of thorax that is smooth and shiny, a large squarish node on the waist, and a large stinger.</p>	<p>MAXFORCE@COMPLETE Brand Granular Insect Bait in stations or broadcast were effective in a field study. DuPont Advion Ant Gel and Optigard Ant Gel Bait performed better than the control in a lab study.</p>	<p>This ant does not form strong foraging trails and does not group recruit.</p>
<p>ANTS - OTHER PB1629</p>	<p>Ants have elbowed antennae, a thread-like waist with one or two bumps. Unmated reproductives have wings, the front wings are larger than the hind wings. Workers ants are wingless. Ants are social insects. Colonies are usually established by a queen. Workers feed the queen, care for the brood and defend the nest. Workers travel along well-marked trails between the nest and food.</p>	<p><u>Baits:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Gourmet Ant Bait Gel Gourmet Ant Bait Liquid Stations Niban FG Optigard Ant Gel Bait PT Advance 375A Select Granular PT 388B Advance Ant Gel Bait Terro-PCO (1%)</p> <p><u>Liquid Baiting Systems:</u> Gourmet Ant Bait Liquid Terro -PCO(1%)</p> <p><u>Pyrethroid Sprays:</u> TalstarP, Tempo SC Ultra Tempo Ultra WP Transport Suspend SC Demand CS</p> <p><u>Slower-acting, non-repellent spray:</u> Termidor SC 0.06% (outdoor perimeter) Phantom SC indoor crack and crevice, outdoor entry points</p>	<p>Follow good sanitary and exclusion practices. Exploit worker caste by using baits. Find foragers and place bait near foraging trail. Workers then bring the poisoned bait back to the nest where it is distributed among all members of the colony.</p> <p>If ants are foraging indoors from an outdoor nest, exclude ants by sealing entry points such as window sills and door steps, or spray entry points into the structure.</p>
<p>BATS PB1624</p>	<p>Night-flying creatures invade attics and similar areas.</p>	<p>Treat area with insecticides to control external parasites including fleas, mites, and bat bugs after bats excluded.</p>	<p>Close entrance holes with one-way excluders to prevent bats from returning after they have left the resting area. Seal opening once all bats excluded. See details in PB 1624.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>BED BUGS PB 1763, PB 1807, SP 761</p>	<p>Flat, oval, reddish, wingless insects. Bloodsucking. Night feeders.</p> <p>Confirm identification. Bat bugs easily confused with bed bugs. Pronotal hairs longer than width of eye for bat bug. If bat bugs are feeding on people, control will not be achieved until bats are excluded.</p>	<p><u>Detection</u> canine detection, especially helpful for inspecting large buildings or areas (check for third party certification)</p> <p><u>Monitoring Devices</u> CLIMBUP Insect interceptor, white or black, http://www.insect-interceptor.com/ (Place bed or furniture leg into center well to aid detection of new infestations and protect against reinfestation. Important to prevent other bed bug access to bed and refill wells with talc as needed.). CLIMBUPS helpful to determine efficacy of treatment when placed additionally in other places around room.</p> <p>BlackOut™ Bed Bug Detector http://blackoutbbdetector.com/. Similar to ClimbUp but does not require retalcing.</p> <p>Many monitors, check for validity by comparing research results.</p> <p><u>Mattress and Boxspring encasements</u> Protect-A-Bed (BugLock™ 3-sided zipper system and ALLERZIP™ seal), Mattress Safe® "Ultimate" zippered encasement Others, ensure it refers to bed bug proofing</p> <p><u>Mattress seams, folds & edges</u> Phantom Aerosol Steri-Fab Bedlam Plus EcoRaider Bed Bug Killer (kills eggs)</p> <p><u>Other cracks and crevices</u> Phantom Aerosol Transport GHP Temprid SC, RTS Tandem Zenprox EC, Aerosol EcoRaider Bed Bug Killer</p> <p><u>Dust for Voids</u> CimeXa Drione PT Tri-Die Pressurized Silica + Pyrethrin Alpine</p> <p><u>Heat treatment (all bed bug stages will die when exposed to 122F)</u> Equipment to heat rooms or buildings, trailers, tents and small chambers is expensive and often requires numerous circuits or generators. For whole room treatments, remote monitoring of temperatures using electronic sensors (minimum 5) AND overhead sprinkler protection that is compliant with fire safety regulations is recommended. A printed report should show that all sensors, including those in tight crevices, reached 122°F. Post-treatment should include an inspecting and use of monitoring devices to determine efficacy, vacuuming dead bed bugs and a perimeter application of DE or silica (see dusts above).</p> <p><u>Chemical Fumigation</u> Vikane</p>	<p>Treat every crack and crevices in bedroom and elsewhere, if needed. Do not spray sheets or blankets. Some PMPs will not treat mattresses with insecticides because of potential human exposure while others will cover treated mattresses and boxsprings with a bedbug proof encasement after treatment. Others may physically remove bed bugs with vacuums and then cover with bed encasement specifically designed to prevent bed bugs from biting through or escaping (i.e., Protect-A-Bed with BugLock™ 3-sided zipper system and ALLERZIP™ seal, or Mattress Safe® "Ultimate" zippered encasement - zipper must be kept closed). Some insecticides may need to contact bed bugs directly to be effective. Resistance to pyrethroid insecticides is widespread. Rotate chemicals used and do not rely on one type (use nonresiduals, residuals and dusts). Foggers are ineffective. Incorporating nonchemical controls including vacuuming, low moisture steaming (AmeriVap, Hi-Tec Cleaning Systems, etc.) laundering (dryer on high for 20 minutes for dry clothes), and removing of infested items may be necessary to manage bed bugs and may be more important as resistant bed bug population are encountered. Cold takes longer to kill bed bugs. Bed bugs need to be exposed to -0 F for 4 days for all stages to die. Reducing clutter will reduce callbacks. Heat or chemical fumigation can be conducted on entire buildings or certain materials, including box springs and mattresses, placed in a permanent or temporary chamber. Fumigation does not provide residual control. Whole heat treatments of rooms should include a perimeter insecticide treatment to prevent bed bugs from moving to surrounding rooms. Wrapping and taping mattresses in black plastic and placing in sun does NOT heat the mattress evenly and does NOT reach the bed bug's lethal thermal threshold on the underside. Difficult to treat items (appliances, lamps, etc.) can be treated with Nuvan Prostrips in bags but may not kill all bed bugs in protected locations. CIRKIL rag-in-a-bag technique (http://cirkil.com/rag) may be more thorough, but odors may linger.</p> <p>See UT Extension publication PB 1763 for more details on bed bug biology. For additional bed bug information and equipment see our <i>Bed Bug in Tennessee Websites</i> at http://bedbugs.utk.edu/</p> <p>The NPMA has released <i>Bed Bug Best Management Practices</i>, http://www.bedbugbmps.org/ so consumers can understand what to expect when working with trained pest management professionals.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>BOOK LICE</p>	<p>Small, soft-bodied, cream-colored to grayish or light brown, wingless, fast-moving. Feed on molds, fungi. Found in books, cereals, wallpaper, boxes. May damage starched things.</p>	<p>PT Microcare Pressurized Pyrethrum Capsule Suspension PT 565 Plus XLO PT 221L PT Cy-Kick CS Pressurized Crack & Crevice Residual PT Tri-Die Silica & Pyrethrum Dust PT Tri-Die Pressurized Silica & Pyrethrin Dust</p>	<p>Large numbers of book lice develop under excessive humid conditions, moldy books, papers, bags or cereals. Dry out infested areas. Destroy infested material of little value. Space sprays of pyrethrins may cause the book lice to scatter throughout structure.</p>
<p>BOXELDER BUGS See SP341-H</p>	<p>Flat, ½ inch long, 1/3 inch wide, dark brownish-black, with 3 lengthwise red stripes behind the head. Wings leathery at base. Membranous at tip with red veins; abdomen is red. Nymphs are smaller, wingless and bright red.</p>	<p>Treat listed sites on label when bugs are first seen. Tempo SC Ultra Tempo Ultra WP Dragnet SFR Demand CS Demon TalstarP Suspend SC Deltadust</p>	<p>These insects are attracted to buildings in the spring and fall. Large numbers collect on siding, around doors, sunny walls and attics. Use exclusion practices before pests become apparent. Inside, vacuum into a dry vac. Avoid use of space sprays if bugs found in wall voids. Dead bugs in wall voids could serve as carpet beetle food. Eliminate female (seed-pod-bearing) boxelder. Outside: vacuum the bugs into water mixed with 1 teaspoon of a liquid detergent per gallon of water in wet/dry vacuum cleaner tank.</p>
<p>BROWN MARMORATED STINK BUG http://eppserver.ag.utk.edu/Whats/wh2009/Issue-1-2009.pdf new web publication expected in 2016</p>	<p>BMSB adults are shield shape, about 5/8 inches long, just about as long as wide, mottled brownish grey with black antennae marked with a whitish band on the next to last segment, dark bands on the membranous part of the wings, and coppery or bluish metallic punctures (small, round depressions) on the head and pronotum). Abdominal segments protruding from the wings are marked with black and white bands. Serious agricultural pest that may overwinter in homes.</p> 	<p>If exclusion methods aren't working completely, they may be supplemented with professionally applied outdoor treatments. Products containing pyrethroids (bifenthrin, beta-cyfluthrin, cyfluthrin and lambda cyhalothrin) and neonicotinoids (acetamiprid and thiomethoxam) have been found to be effective against brown marmorated stink bug. In general, pyrethroids are faster acting than other chemistries; however, new pesticide labels limit professionally applied pyrethroids to 1 inch bands around windows and doors when the surface is over a hardscape. Insecticides will have limited persistence outdoors in the sunlight and rain and may not prevent the brown marmorated stink bug from entering structures.</p> <p>Alpine WSG Talstar P Suspend Polyzone Other pyrethroids (check label)</p>	<p>Pest-Proof by late summer using techniques described at https://ag.tennessee.edu/EPP/Redbook/53%20modhomepestvailmod22615a.pdf. If supplemental pesticide applications are deemed necessary, make exterior spot, crack & crevice and/or void applications where these pests may harbor or hibernate, such as cracks and crevices, in weep holes, wall voids, around window and door frames, attic vents and behind siding, in late summer/early fall before the pests arrive. Pyrethroid labels are more restrictive so read label carefully. Indoor light traps may help reduce populations inside. Vacuum bugs found inside into knee-hi stocking placed on the end of the vacuum tube prior to attachment placement. After vacuuming, remove knee-hi, tie off, place in a sealed bag and in outside garbage can. If BMSB found on walls in large numbers, they can be removed by using an empty plastic milk jug with the bottom removed. Invert the jug, ensure the cap is tightly closed and add a small amount of soapy water. Likewise, the top of a capless plastic soda bottle can be removed at the widest part of the neck, inverted and placed back on the bottle to create a funnel trap. When the edge of the modified jug or bottle is moved up the wall towards a BMSB aggregation, the bugs will drop into the trap or soapy water. Soda bottles can be sealed in a plastic bag and placed in the freezer for a few days. Frozen or drown bugs can be disposed of outdoors in a garbage can or compost pile. Flushing BMSB down the toilet will waste water and is not recommended.</p>


PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>CARPENTER BEES SP341-P</p>	<p>½ to 4/5 inch long with a blue-black, green or purple metallic sheen. Color and size resembles a bumble bee, but the top of the abdomen is hairless.</p> <p>These bees chew a circular, 3/8-inch entrance hole into wood and nesting gallery 4-6 inches long at a right angle to the entrance hole. Galleries used for several years may extend 10 feet.</p> <p>Carpenter bees overwinter in previously used galleries, so re-inspect and seal galleries in the fall.</p>	<p>Apply dusts into the gallery openings: Tempo 1D DeltaDust Apicide</p> <p><u>Sprays:</u> Tempo SC Ultra (spray or foam) Tempo Ultra WP (spray or foam) PT Cy-Kick Dragnet SFR Suspend SC 0.06% Premise .05%-0.1% (spray or foam)</p>	<p>Use a badminton racket to kill flying adults or use a straight wire to break up cells in tunnels. Individual bees can be caught with a net and killed.</p> <p>In the spring, apply foam, spray or puff of insecticidal dust into nest holes in the evening when the carpenter bees are at rest. Allow bees access to the nest for 24 hours. Seal the hole with putty, 3/8 inch diameter dowel or cork to prevent reinfestation.</p> <p>Dusts are the preferred formulation because they coat the porous wood surface of the bee's gallery.</p>
<p>CARPET BEETLES</p> <p>Black Carpet Beetle</p> <p>See SP341-I</p> <p>Common; or Furniture; or Varied carpet Beetles</p>	<p>Adult 1/8 inch to 1/4 inch in length; black; brown legs. Larva 3/8 inch in length; carrot-shaped; covered with short hair and has long terminal bristles.</p> <p>Adults 1/8 inch long with white and orange; or yellow, white and black; or white, brown and yellow spots; larva with long black to brown hairs.</p>	<p>Treat cracks, crevices and hidden area of walls, closets, stored materials, under carpets, etc. Do not apply insecticide to clothing. See precautionary statements about pesticides staining carpets.</p> <p>Tempo Ultra WP Tempo SC Ultra PT Cy-Kick Demand CS Dragnet SFR Suspend SC DeltaDust Kicker PT Tri-Die Silica & Pyrethrum Dust PT 565 Plus XLO PT Microcare CS Controlled Release Pyrethrum PT Microcare Pressurized Pyrethrum Capsule Suspension</p>	<p>Carpet beetles infest carpeting, clothing, fur, upholstered furniture, books, bird nests, milk powders, articles of animal products, feathers, wool, silk and other materials of animal origin.</p> <p>Locate food source and remove, if possible. Use good housekeeping such as cleaning floor and carpets regularly. Dry clean clothes regularly. Stored materials subject to damage should be thoroughly cleaned and stored in tight container with moth crystals.</p>
<p>CENTIPEDES</p>	<p>Grayish creature with long feelers and many long, slim legs. Fast moving. Long antennae.</p>	<p><u>Indoors:</u> DeltaDust PT Tri-Die Pressurized Silica & Pyrethrin Dust PT Tri-Die Silica & Pyrethrum Dust Talstar P PT Cy-Kick Crack & Crevice Pressurized Residual PT Cy-Kick CS Controlled Release Cyfluthrin Tempo SC Ultra PT Microcare CS Controlled Release Pyrethrum</p> <p><u>Outside Perimeter:</u> Demon WP DeltaDust Tempo Ultra WP Tempo SC Ultra Cy-Kick TalstarP Talstar granulars DeltaGard G</p>	<p>Feed on insects. Can bite. Usually not numerous. Active at night. Non-chemical control: leave a 12-18 inch bare zone next to foundation base. Move wood piles and other clutter away from structure. Spot-treat cracks and crevices, door thresholds and moldings where pest may crawl or as directed by label. Residuals may not provide total control. Treatments may need to come into direct contact with pest.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
CHIGGERS	Very tiny mites which get on the person and cause blotches and itching. Don't usually invade homes, but occur in yards and wooded areas.	Use commercial repellents around ankles and waist for personal protection. Apply deet repellent to skin; and Permanone 0.5% spray to shoes, cuffs and socks, heed drying directions before wearing. Apply to infested areas of turf. Tempo SC Ultra Tempo Ultra WP Talstar P	Mow lawn regularly. Remove weeds and brambles from fence rows.
CLOTHES MOTHS SP341-J	Brownish moths, wingspread 1/2" long. Larvae are 1/16-1/3" long. Gray silken cases or webbing over surface of fabric.	Vacuum prior to treatment and to remove potential food sources. Apply sprays to cracks and crevices in closets and spot treat other infested areas. Tempo Ultra WP Tempo SC Ultra Cy-Kick DeltaDust Suspend SC Kicker	Adults are not attracted to light and will fly to dark corner when disturbed. Larvae usually found on infested materials, wool, fur, feathers, hair, upholstered furniture, nonsynthetic carpets, dust and lint. Do not spray clothes. Store susceptible items with moth crystals in sealed storage container. Heed warning about staining clothing.
CLOVER MITES	Tiny (1/30 inch) mites, brown to olive green with pair of long front legs.	Apply sprays to points of entry such as foundations, windows and doors. Cy-Kick CS PT 221L TalstarP Mavrik Perimeter	Invade homes from the yard in great numbers in fall and spring. Stain walls or fabrics reddish-brown when crushed. Keep grass and shrubs from direct contact with house. Good weed control in turf and a vegetation free border of 12-18 inches around home will help.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>COCKROACH PB1024</p> <p>German Cockroach</p> <p>Brown Banded or Furniture Cockroach</p> <p>Oriental Cockroach</p> <p>American Cockroach</p> <p>Smokybrown Cockroach</p>	<p>About 5/8" in length, pale brown or tan with 2 parallel dark streaks on pronotum. Usually most abundant in the kitchen and bathrooms.</p> <p>1/2" to 5/8" in length, dark brown with 2 pale bands traversing wings. Widely distributed throughout the house in walls, closets, furnishing, in appliances, but abundant in kitchens.</p> <p>1 1/4" in length, dark reddish-brown to black, wings do not surpass end of abdomen. Usually found in lower floors, outdoors or in crawl space. Frequents water meters, floor drains or moist, dark areas.</p> <p>1 1/2" in length, reddish-brown with pale yellow band around pronotum. May be found throughout house, outdoors, in crawl spaces, sewers, water meters and garbage cans.</p> <p>1 to 1 1/2 inches, uniform, very dark brown to black; head shield is a solid, dark color. Takes harborage in moist, warm and dark places like tree holes, mulches, soffits in attics with poor ventilation.</p>	<p>Do not spray repellent insecticides near baits.</p> <p><u>Baits in cracks and crevices:</u> Dupont Advion Cockroach Gel Bait Dupont Advion Cockroach Bait Arena Maxforce Roach Killer Small Bait Stations Maxforce Roach Killer Bait Gel Maxforce FC Select Roach Killer Bait Gel (for bait averse roaches) Maxforce Impact Roach Gel Bait Maxforce FC Magnum Roach Killer Bait Gel Niban G Prescription Treatment Advance Cockroach Gel Bait Reservoir Prescription Treatment Avert Dry Flowable Cockroach Bait Form. 1 Vendetta</p> <p><u>Baits for large roaches:</u> Dupont Advion Cockroach Gel Bait Maxforce Granular Bait Maxforce Roach Killer Bait Gel Maxforce FC Magnum Roach Killer Bait Gel MAXFORCE@COMPLETE Brand Granular Insect Bait Maxforce Impact Roach Gel Bait Niban G Vendetta</p> <p><u>Insect Growth Regulators for small roaches</u> containing hydroprene (Gentrol Aerosol, Gentrol IGR Concentrate, Gentrol Point Source) or pyriproxyfen (Archer, Nylar or others). IGRs are slower acting but longer lasting - sterilizes adults.</p> <p><u>Lightly dust voids with:</u> PT Tri-Die Drione DeltaDust NiBor-D</p> <p><u>Crack and Crevice Sprays:</u> TempoUltra WP, SC Ultra PT Cy-Kick CS Pressurized Crack & Crevice Residual Suspend Talstar P PT 221L Phantom</p> <p><u>Outdoor perimeter for large roaches</u> PT Cy-Kick CS Controlled Release Cyfluthrin Suspend TempoUltra WP DeltaGard G Niban FG, G MaxForce Granular Bait</p>	<p>Prevent access to food, water and shelter. Practice good sanitation in food handling, storage and eating areas. Control moisture, prevent leaks or condensation. Seal off harborage sites such as cracks and crevices with caulk, etc. Also use exclusion practices to prevent cockroach movement.</p> <p>Use glue boards or sticky traps placed along edges in dark places to locate and monitor cockroach populations.</p> <p>Baiting is the preferred method for cockroach control. Apply baits to cracks, crevices, pipe opening into walls, joints of furniture and cabinets, pipe conduits, and elsewhere as indicated by glue board catches.</p> <p>If you chose to spray, use precautions to keep chemicals out of food, spices, and off dishes or eating utensils. Do not apply sprays where electrical shorts may occur; use baits or dusts in these areas.</p> <p>Do not use sprays when baiting because cockroaches may be repelled from the baits. Read label carefully; some products may not be labeled for food handling areas.</p> <p>Increase ventilation in attic to reduce attraction to smokybrowns.</p> <p>American cockroaches often enter facilities through drains with a dry p-trap. Items, such as ProSet Tap Guard or Sure Seal Inline Floor Drain, prevent sewer gasses and cockroaches from escaping into living spaces, but still allow water to drain.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>CRICKETS</p>	<p>Black, jumping insects with long antennae. Cave or camel crickets are humped and brown.</p>	<p><u>Baits:</u> Maxforce Granular Bait Niban Granular Bait</p> <p><u>Sprays and dusts:</u> Tempo Ultra WP Tempo SC Ultra Cy-Kick Demon WP Demand CS Talstar P DeltaDust Suspend SC</p>	<p>Black cricket may damage clothing. Repeat treatment as needed. Spray entry points into structure. Dust crawl space.</p> <p>Camel crickets infest damp basements, under slabs and crawl spaces. Ventilate or dry these areas. Active at night. Apply sprays into cracks and crevices where crickets dwell.</p> <p>Use exclusion practices. Glue boards can be used indoors around entry points and other places in basements, etc.</p>
<p>EARWIGS</p>	<p>Easily identified by pair of "pinchers" at end of abdomen.</p>	<p><u>Outdoors:</u> Talstar P Demon W Dagnet SFR Demand CS PT Cy-Kick CS Controlled Release Cyfluthrin</p>	<p>Earwigs are incidental invaders into houses. They usually dwell in leaf litter, mulch and woodpiles and are common "hitchhikers" on vegetables harvested from the garden.</p> <p>Moving compost piles away from the house will aid pest control. Insecticidal control is usually unnecessary for this insect.</p> <p>If needed, spray possible entry points and mulched area around the house.</p>
<p>FLEAS PB 1596</p>	<p>Small, 1/16" long, reddish-brown, wingless insect. Body compressed laterally, legs long and adapted for jumping.</p>	<p>On pet:</p> <p>CATS, KITTENS, & PUPPIES ARE MORE SENSITIVE TO INSECTICIDES!!!! Consult a veterinarian and always read label prior to treating a pet.</p> <p>Veterinarian supplied products - usually kill fleas within 12 - 36 h or sooner and provide 90 - 95% control for about 30 days: see pesticide recommendation at http://www.ent.uga.edu/pmh/ under <i>Homeowner</i>, and <i>Animals</i>, for a thorough listing of veterinarian supplied on-pet products.</p> <p>Indoor Premise C & C Treatment Spot-treat infested areas and pet resting areas inside with following: Insect Growth Regulators [and adulticides]: pyriproxyfen (Archer and others) pyriproxyfen [and permethrin or pyrethrin] (PT Ultracide and other ready-to-use products) methoprene (Precor IGR Concentrate) methoprene [and pyrethrins or permethrin] (Precor 2000 Premise Plus Spray and other ready-to-use products)</p> <p><u>Adulticides</u> Suspend SC, Alpine WSG, others</p> <p>Outdoors (when specified on label): pyriproxyfen (Archer and others); Alpine WSG Demand CS (outdoors only); Suspend Talstar P Talstar PL Granular</p>	<p>Keep pets and people out of treated area (indoors and outdoors) until spray dries.</p> <p>Step 1. With veterinarian supplied products that are currently available, control of fleas in small- to moderate-sized infestations is likely to occur by using those pet treatment products alone. May take 2 months to completely break flea life cycle.</p> <p>Sprays of pyrethrum and pyrethroids may not work as well as the newer chemistry used in pet treatments because of insecticide resistance.</p> <p>If pet treatment alone does not provide sufficient control, initiate a complete control program by April.</p> <p>Step 2. Vacuum infested areas twice a week and prior to treatment to remove eggs, larvae, adults and organic matter. Steam- cleaning carpet may also reduce populations. Eliminate fleas from pets, bedding and premises before departing on vacation.</p> <p>Step 3. Treat pet resting areas indoors and clean or remove pet bedding on the same day. Insect growth regulators important to break flea life cycle. A combination of an insect growth regulator and an adulticide may be the most efficient formulation to use.</p> <p>Step 4. Mow grass, keep weeds down and trim shrubs to expose flea eggs and larvae to lethal dessication. Irrigating areas surrounding buildings, but not against building, may kill fleas by drowning. If fleas are surviving outdoors, apply insecticide to labeled areas.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>FLIES Face flies, cluster flies, and blue bottle flies</p>	<p>Adult flies of these three species hibernate in attics and wall voids. Cluster flies about 1/3" long, dark gray, with checkered black and silver abdomen, with gold hairs on thorax of newly emerged adults. Face fly similar in appearance to the house fly. Adult blue bottle flies have a dull gray thorax and a shiny blue abdomen.</p>	<p>Exclude flies in the fall by sealing entry points, screening behind all vents, sealing holes in walls and attics prior to pest entry in fall.</p> <p>Can apply pyrethroids to potential entry points prior to pest entry. Vacuum or use pyrethrin sprays to kill exposed flies. Use black light trap with sticky surface.</p> <p>Dust voids (boric acid not very effective) where flies may be over-wintering.</p>	<p>Cluster fly larvae are parasites of earth worms.</p> <p>Face fly larvae develop in cow patties. Adults overwinter in attics and wall voids.</p> <p>Blue bottle fly larvae develop in garbage, decaying meat, dead animals, fish and excrement.</p>
<p>House flies</p>	<p>About 1/4" in length, dull gray color with 4 longitudinal dark stripes on the thorax.</p>	<p>Remove larval food sources.</p> <p><u>Spray outdoor areas where flies rest such as garbage collection sites:</u> Tempo Ultra WP, Demon WP Demand CS, Dagnet SC, Suspend SC, CyKick CS, Cynoff</p> <p><u>Baits (rotate use of diamide, carbamate, and neonicotinoid):</u> For use around commercial facilities. Should not be used inside or around homes, or any other place where children or pets are likely to be present. Zyrox Fly Granular Bait (diamide) Golden Malrin Fly Bait (carbamate) Maxforce Granular Fly Bait (neonicotinoid) Maxforce Fly Spot Bait (neonicotinoid) Florida-Fly Baiter with Maxforce Fly Spot Bait (neonicotinoid)</p> <p><u>EndZone Insecticide Sticker</u> Stickers work best when placed on or near a window or other light source. However, in the absence of light or under low-light conditions, stickers may be placed near a potential fly food source (such as inside a garbage can).</p> <p>If needed, use pyrethrin space spray for temporary relief inside home.</p>	<p>Larvae develop in warm organic matter of animal or vegetable origin. Remove trash at least twice a week to reduce fly populations in homes. Screen windows and doors. Garbage cans should have tight-fitting lids. Use insect light traps indoors. Sanitation is very important.</p>
<p><u>Bottle Flies</u> Green Bottle Fly Blue Bottle Fly Bronze Bottle Fly Black Bottle Fly</p>	<p>1/2" in length; green metallic color 1/2" in length; blue metallic color 1/2" in length; bronze metallic color 1/3" in length; shiny grey thorax and dull blue metallic abdomen.</p>	<p>Remove larval food sources.</p> <p>Spray outdoor areas where flies rest or try an insecticidal sticker (EndZone) indoors to reduce adult populations.</p>	<p>Bottle flies indoors often indicate a dead mouse or other animal in wall voids, attic, basement, etc.</p> <p>Dispose of dead animal carcasses, animal excrement, and other potential larval food sources such as decaying vegetation and garbage.</p> <p>Garbage cans should have tight-fitting lids.</p>
<p>Small Fruit Fly or Vinegar Fly</p>	<p>1/8" long, red eyes, tan head and thorax, abdomen gray-black. Some species have darker eyes, see http://eppserver.ag.utk.edu/Whats/wh2006/Issue-11-2006.htm</p>	<p>Remove larval food sources. Check mops and brooms too. Use a wire brush, or foam or other application of microbials or botanical drain cleaners to labeled sites (which may include drains, baseboards, behind bars, drip trays, bundles of sticky syrup lines, grout ruts, under ice machines and other equipment), that should be cleaned, but are often neglected and difficult to clean. May need to be repeated. Traps may reduce adult fly populations. Pyrethrins for temporary relief of adults.</p>	<p>Egg to adult in 8-11 days. Larvae in decaying fruit, vegetables and garbage cans, etc. Adults around larvae.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
Moth Fly Sewer Fly Drain Fly	Small, scaly or hairy, long-legged moth-like flies. Wing veins parallel.	Remove larval food sources. Remove moist organic materials, clean drains with wire brush or steam clean. Following initial mechanical cleaning in commercial accounts, botanical, microbial and/or enzymatic drain cleaners can be applied to labeled sites, which may include drains, waste water traps, etc. to maintain clean surfaces. A foam formulation may work best on vertical surfaces. Pyrethrins for temporary relief of adults.	Adults rest on walls or foliage. 3-4 weeks from egg to adult. Larvae found in slimy drains, sewer backup or leaks, unsanitary garbage cans, potted plant saucers, baths or feeders for birds, clogged roof gutters or storm drains, drip lines from air conditioners, moist compost, septic tanks and other places that hold very moist organic solids.
Phorid, Humpbacked Fly, or Scuttle Fly	Adult small, 1/16 - 1/8" long; brown, black or yellow; thorax humped when viewed from side. Dark veins along front edge of wing. Adult scuttles or "runs" erratically over surface.	Remove larval food sources. Check bottom of trash cans, cracks under appliances/equipment, garbage disposals, rotting vegetables and meats, mop heads, septic systems, and potted plants that have been overwatered, flowers in vases, mausoleums, and soiled bedding of animals. Use insect light traps to catch adults to determine if potential larval source nearby and to temporarily reduce adult populations. If source cannot be found, consider a cracked sewer or waste pipe. If sewer pipe the cause, remove slab, repair pipe and remove contaminated soil. Insecticidal sticker (EndZone) may also help reduce adult fly population.	Development time 11 days (85F) to 28 days (72F) for common species. Larvae found in moist decaying organic matter (feces, carrion, fungi and decaying plants). Phorid flies were abundant in 2009 and sometimes the larval source was outside in decaying vegetation near home.
FUNGUS GNATS SP341-C	Adults 1/8 to 1/4 inch long. Slender larvae have shiny black head and white thread-like body. Adults attracted to light. Collect in windows and soil in potted plants. Run rapidly over surface.	Pyrethrins, insecticidal sticker (EndZone) or light traps may also help reduce adult fly population caused by moisture and decay from leaks including roofs. See SP 341-C for list of products labeled for adult and larval fungus gnats in interiorscapes. Need a category 3 certification and working under someone licensed in HRI to make interiorscape pesticide applications.	Avoid over-watering plants, let soil dry out somewhat between waterings so larvae don't have fungi to feed on. If plants not the problem, look for water leaks or other moisture problems. Check flat roof and piles of pet bird droppings which can support fungal growth. Remove moisture or dry moist areas. May enter from outdoors. Rake and reduce mulch depth to 2 - 3 inches.
HEAD LICE SP341-S	Tiny, flat insects which infest people and clothing.	Premise sprays are not recommended for head lice control. Several louse shampoos and other hair products are available for homeowner use.	Wash infested clothing and bedding with strong soap and very hot water; tumble dry on high heat. Dry clean woolens. Do not share hair brushes, caps, etc. Use special combs to remove nits (eggs). Nits hatch by 10 days, so another application of head lice shampoo may be needed at this time. Follow label directions.
KUDZU BUG	¼-inch long, almost square-shaped with a brown to olive-green color. The immature stages are similarly shaped but smaller and "hairy."  A map of the state of Tennessee showing the distribution of Kudzu Bugs from 2012 to 2015. The map is color-coded by county: blue for 2012, green for 2013, yellow for 2014, and red for 2015. The distribution shows a general trend from the western and northern parts of the state towards the eastern and southern parts over the four-year period. A legend in the bottom right corner identifies the colors for each year. A note at the bottom left of the map reads: "Kudzu Bug Distribution 2012-2015" and "©2016 University of Tennessee System".	If exclusion methods aren't working completely, they may be supplemented with professionally applied outdoor treatments. Products containing indoxacarb, dinotefuran, pyrethroids (such as b-cyfluthrin, bifenthrin, cyfluthrin, deltamethrin and λ-cyhalothrin), or pyrethroids combined with neonicotinoids (imidacloprid, acetamiprd, or thiomathoxam) have been found to be effective against kudzu bug. Insecticides should be applied around windows, doors and other entry points as is done for other occasional invaders. In general, pyrethroids are faster acting than other chemistries; however, new pesticide labels limit professionally applied pyrethroids to 1 inch bands around windows and doors when the surface is over a hardscape. Alpine WSG Tandem Talstar P Other pyrethroids and combination neonicotinoid & pyrethroids (check label)	Exclude the pests before they start aggregating on structures to overwinter. Make spot, Crack & Crevice and/or void applications where these pests may harbor or hibernate, such as cracks and crevices, in weep holes, wall voids, around window and door frames, attics and behind siding. Apply to exterior wall surfaces around entry points and resting areas where insects congregate. Vacuum bugs found inside into knee-hi stocking placed on the end of the vacuum tube prior to attachment placement. After vacuuming, remove knee-hi, tie off, and dispose in soapy water. Insecticides will have limited persistence outdoors in the sunlight and rain and may not prevent kudzu bugs from entering structures.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
LADY BEETLE, MULTICOLOR ASIAN SP503-C	Multicolored Asian Lady Beetles (MALB) start searching for overwintering sites, your home, on the first or second day when temperatures are greater the 65°F after a dramatic drop in temperature, usually to near freezing. This usually occurs about the third week in October.	Demand CS Suspend SC TalstarP other pyrethroids	1. Pest-Proof: seal entry points before beetles arrive. 2. Treat roof lines or soffits, vertical contrast areas, and entry points (around the following outdoor items: windows, doors, vents, pipe penetrations) with pesticides before the beetles arrive. 3. Remove dead beetles as they pile up because they may cause other MALB to aggregate. 4. If the beetles make their way into the home, vacuum (with HEPA to prevent allergen circulation) or try a light trap.
MICE PB1624	Adults weigh about ½ ounce. Dusky gray color, slender body, prominent ears, tail about as long as head and body.	Place snap traps, multiple catch traps and glue boards along paths traveled by mice. Traps or glue boards should be placed every 8-12 ft. Traps can be baited with: dry rolled oatmeal, bacon squares, small wads of cotton or gumdrops. Baited traps should be set at right angles to rodent runs. Place trap at right angles to rodent pathway with trigger part of trap against the run. EPA has changed allowances regarding use of rodenticide baits in the urban environment. Check for latest updates before using baits. http://www.epa.gov/oppsrrd1/reregistration/rodenticides/finalriskdecision.htm	Mice move in from outdoors in fall as temperatures decline. Exclusion practices needed, mice can fit through an opening 1/4" in diameter. Sanitation: remove access to food, water and shelter. Rodents use edges of walls, studs and pipes as guidelines. Remember to set traps where children and pets will not be hurt. Mice are curious and will normally approach traps the first night. If you don't catch a mouse in the first few nights, the trap is in the wrong location.
MILLIPEDES	Slender, brownish, multi-legged, hard-shelled, 1-2" long. Two pair of legs per body segment. Invade home from outdoors. Harmless.	<u>Outdoors:</u> Cy-Kick CS Cynoff WP Demand CS Suspend SC Talstar P Talstar PL (granular) Tempo SC Ultra Tempo Ultra WP	Millipedes are not insects, so insecticides are not always effective. Best control obtained when pest comes in direct contact with the insecticide. Usually occasional invaders and may invade in large numbers. Under these circumstances, non-chemical control may be more effective: remove mulch and other clutter from near the foundation, dethatch lawns and water in the morning. Prune tree limbs to dry their habitat. Use exclusion practices. Treat entry points into structure, shady areas, ivy beds, flower beds and rock walls, leaf-litter or as directed by label. Millipedes will die within 2-3 days after entering a dry structure.
MITES, BIRD OR RODENT	Mites occasionally found indoors because of rodent or bird nest in, on, or near structures. Some of these mites may bite people. They are small (about the size of a period), but can usually be seen with the naked eye.	Dust or spray cracks and crevices around infested area. Tempo 1D (as for other pests) DeltaDust (as for other pests) Talstar P (as for other pests) If widespread, space sprays of pyrethrins may be necessary.	The first step in controlling bird or rodent mites is to eliminate the host animals and remove their nesting sites. Often, the nests will be found in the attic, around the eaves and rafters or in the gutters or chimney. Gloves should be used when handling dead animals. A respirator should also be worn when removing nest materials to avoid inhaling fungal spores and other potential disease-producing organisms associated with the droppings. See chiggers for repellents.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>MOSQUITOES SP503-B</p>	<p>Delicate insects that bite humans and animals. Larvae and pupae in water. Adults stay in shrubbery, crawl spaces, etc.</p> <p>Discharges to waters of the U.S. from the application of pesticides will require NPDES permits starting April 9, 2011 (http://cfpub.epa.gov/npdes/home.cfm?program_id=410) and applicators will need to abide by the state of Tennessee's National Pollutant Discharge Elimination System (NPDES) permit. The state NPDES permit will be issued through the Tennessee Department of Environment and Conservation at http://state.tn.us/environment/</p> <p>See the new mosquito control association, Tennessee Mosquito and Vector Control Association, http://tennmosquito.org, for training opportunities.</p>	<p><u>Treat standing water with:</u> <i>Bacillus thuringiensis israeliensis</i> (Bti):</p> <p>Aquabac xt, 200G Teknar HP-D and G, Vectobac G, GS and 12AS</p> <p>methoprene (IGR) Altosid SBG, XR-G, Pellets, Liquid, XR- briquets, briquets, Pro-G Pre-Strike Granules</p> <p>Spinosad Natular 2E, G, G30, XRT, T30</p> <p>temephos Abate Pellets Abate Tire Treatment</p> <p><u>For <i>Culex</i> mosquitoes in septic conditions, treat standing water with:</u> <i>Bacillus sphaericus</i> Vectolex CG, WDG, WSP</p> <p><u>Outside Area ULV applications by ground equipment for use by trained professional personnel:</u> Anvil 10 + 10 Biomist 3 + 15 ULV Biomist 1.5 + 7.5 Biomist 30 + 30 ULV Kontrol 30 + 30 Mosquitomist One Scourge 1.5 + 4.5 (RUP)</p> <p><u>Outside residual barrier</u> (permethrin, malathion, Tempo, Suspend, Talstar and others) can be applied to vegetation on perimeter of property that is prone to rapid infestation of mosquitoes. This kills adults feeding on nectar or resting in these sites and some may act as a repellent. <i>Aedes</i> species found closer to ground (<10 ft) and <i>Culex</i> found higher in the canopy. Spray other shady damp areas where mosquitoes rest.</p> <p>If needed indoors, use sprays containing pyrethrins in closets, stairwells, behind and beneath furniture for temporary relief.</p> <p>A more extensive list of mosquito control products for commercial and government agencies can be found at</p> <p>http://www.ent.uga.edu/pmh/Com_Humans.pdf</p> <p>These products have not been verified to be registered in Tennessee.</p>	<p>To reduce mosquito populations, a complete mosquito control program must be followed. See UT Extension publication PB 1685 The Tennessee Mosquito Control Handbook (Pesticide Applicator Licensing Manual) for more details.</p> <p>Pesticide applicators applying mosquito control pesticides on public lands and waters need to be certified in category 8 and licensed in PHMC.</p> <p>Eliminate larval sites (standing water) around structure by unclogging gutters, emptying bird baths, children's pools, pet bowls, flower pot saucers, old tires, and other containers around home twice a week. Drain or fill low areas where water collects. Easiest to control mosquitoes in immature stage because confined to water. Treat standing water with labeled insecticide. Repair screens.</p> <p>People should wear repellents when potentially exposed to mosquitoes. Apply N,N-diethyl-m-toluamide or N,N-diethyl-3-methylbenamide called DEET to skin. American Academy of Pediatrics (2003): concentration of 10% DEET appears to be as safe as products with a conc. of 30% when used according to label. Most guidelines cite that it is acceptable to use repellents containing DEET on children over 2 years of age. Children and adults can wear clothing with long pants and long sleeves while outdoors. DEET or other repellents such as permethrin can also be applied to clothing (don't use permethrin on skin), as mosquitoes may bite through thin fabric.</p> <p>On April 22, 2005 the CDC, Center for Disease Control and Prevention, added two new active ingredients, picaridin and oil of lemon eucalyptus, as suggested repellents to prevent mosquito bites.</p> <p>Mosquito netting can be used over infant carriers. More information on repellents and their safe use can be found at http://www.cdc.gov/ncidod/dvbid/westnile/qa/insect_repellent.htm</p> <p>Reduce the number of areas where adult mosquitoes can find shelter by cutting down weeds adjacent to the foundation and in yards, and mowing the lawn regularly.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>PANTRY PESTS</p> <p>Saw-Toothed Grain Beetle</p> <p>Cigarette Beetle</p> <p>Indianmeal Moth</p> <p>Rice Weevil</p> <p>Confused Flour Beetle</p> <p>Red Flour Beetle</p>	<p>Brownish black, 1/8" long, flattened with 6 saw-tooth like projections on thorax. Feeds in a wide variety of stored products, cereals, nuts, dried fruit, cookies, candy, etc.</p> <p>1/8" long, oval, reddish brown, head not visible from above, antennae saw-like. General feeder in tobacco, seasonings especially paprika, cereal, dried flowers, and a wide variety of stored foods.</p> <p>3/8" wing span, inner 2/3 of wing grayish, outer 1/3 of wing mottled copper and black. Feeds in coarse grain products, chocolate, nuts, dried fruit.</p> <p>1/8" long. Long snout on head, dark brown with 4 bright spots on wing cover. Feeds in grains.</p> <p>1/8" long, reddish-brown, antennae gradually enlarged to end in a club. Cannot fly. Feeds in flour and cereal products.</p> <p>1/8" long, reddish-brown, antennae has a distinct 3-segmented club and can fly. Feeds in flour and cereal products.</p>	<p>Locate food source and discard.</p> <p>Place grains, flours, nuts and other stored products in insect proof containers when they are brought home from store.</p> <p>Pheromone traps can indicate the presence of pests and may provide control without insecticides when populations are low and pests confined.</p> <p>Vacuum cracks and crevices and wipe down pantry to remove pests and food source. Do this before insecticidal application, also.</p> <p>Do not contaminate food, dishes, shelves or utensils with pesticides. If insecticide application desired, direct spray into cracks and crevices of storage cabinet shelves when shelves are clean and all food and utensils, etc. removed.</p> <p>Treat only cracks and crevices in the pantry. Do not wash off insecticide residue. Cover with paper if desired.</p> <p>pyrethrins Tempo SC Ultra Cy-Kick DeltaDust Suspend SC PT Cy-Kick CS Pressurized Crack & Crevise Residual PT Microcare Pressurized Pyrethrum Capsule Suspension PT Tri-Die Pressurized Silica & Pyrethrin Dust Drione</p>	<p>To prevent infestations:</p> <ol style="list-style-type: none"> 1) Inspect stored products periodically, 2) practice good sanitation, 3) rotate stored product use so older stores are used first and none remain in storage indefinitely, 4) have adequate ventilation to prevent moisture buildup in storage areas. 5) Insect proofing; use insect-proof package or storage procedures wherever possible. 6) Pheromone traps can indicate the presence of pests and are available for: Indianmeal moth, saw-toothed grain beetle, confused and red flour beetle, cigarette beetle, drugstore beetle, clothes moths and others. <p>Non-chemical control:</p> <p>Either destroy the infested products or salvage them by super heating to 140 degrees F for ½ hour, or super cooling in a deep freeze at 0 degree F for 4 days.</p> <p>Store insect-free beans in containers with tight lids.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>POWDERPOST AND OTHER WOOD-BORING BEETLES E&PP #391</p> <p>Powderpost Beetles</p> <p>Lyctid powderpost beetle</p> <p>Anobiid powderpost beetle</p> <p>Roundheaded borers</p> <p>Old house borers</p> <p>Others</p>	<p>Shot-sized holes along with flour-like powder indicate these beetles.</p> <p>Attacks hardwoods such as oak, ash and hickory found in solid and laminate ring porous hardwood floors and furniture; molding, window and door frames, and wood paneling. Antennae with 2-segmented club. Frass smooth, not gritty. Head protrudes forward. Reinfests seasoned wood less than 7 - 10 yrs. old.</p> <p>Attack hardwoods and softwoods. In addition to above, they also attack beams, rafters, joists, studs and other structural framing. Infestations found in moist, poorly ventilated areas such as crawl spaces, basements, etc. Frass gritty. Head hidden by pronotum. Reinfest seasoned wood that may be decades old.</p> <p>Presence indicated by large hard-shelled beetles with long antennae.</p> <p>Broadly-oval 1/4" emergence hole made by old house borer. Larvae in tunnels packed with frass; 3 eye spots to left and right of mandibles. Beetle 3/4 inch long, grey-brown with 2 patches on wing covers; 2 bumps on thorax. Reinfests seasoned softwoods (pine).</p> <p>Neat 1/2" holes may appear in walls where beetles emerge. Don't usually reinfest seasoned softwoods (pine).</p>	<p>Products listed in this column refer to treatment for all reinfesting wood-boring beetles listed.</p> <p>Beetles that have pupated prior to insecticide application may be unaffected and may continue to emerge. Insecticide applications should prevent reinfestation. Products containing disodium octaborate tetrahydrate (DOT) and glycols (Bora-care, etc.), may penetrate wood further than other residuals, but penetration is variable and depends on moisture content of the wood and other factors. Other DOT products include Tim-bor, Armor-Guard and others. DOT treatments must be made to unfinished surfaces as they will not penetrate paint or varnish; sand or power wash logs prior to treatment. Paint, spray, inject or brush on.</p> <p>If infestation spreads into walls or between floors, fumigation may be needed. Fumigation is costly and should only be considered as a last resort. If only small articles infested such as furniture, antiques, etc., they can be fumigated in a chamber at a lower cost. Only professional pest control operators licensed to fumigate can perform this operation.</p> <p><i>0080-06-14-.08 When it is determined that an active infestation exists, treatment will be permitted for the control or prevention of reinfestation of the families of beetles which are known to reinfest seasoned wood... Preventive treatment in the absence of an infestation is prohibited. ...When wood destroying beetles are present at or below the subfloor level, then control measures should be applied from underneath the structure using any approved pesticide labeled for that use. If there is evidence to indicate or reasonable cause to suspect that a substantial active infestation of wood destroying beetles exists above the subfloor level, then fumigation with an approved fumigant is permitted..... (This rule is expected to change in 2016.)</i></p>	<p>Determine extent of infestation. Signs for powder post beetles are: flour-like "frass" dropping from pinhead-sized or slight larger holes. Anobiids' frass are more gritty than Lyctids; adult beetles attracted to light may be found on window sills or foundation vents. Important to determine if infestation active or not. Mark or seal existing holes, vacuum existing sawdust, recheck wood for new holes in spring or early summer. These beetles damage wood slowly. If "frass" is yellow, caked or covered with dust or debris, that damage is old. Old house borers can be detected by hollow sound when wood tapped.</p> <p>Prevention:</p> <ol style="list-style-type: none"> 1) don't use old lumber from a barn or wood pile unless it has been treated (2) don't use improperly dried or stored lumber (3)inspect firewood prior to bringing into structure (4) paint, varnish or otherwise seal wood to prevent exposed edges (5) seal previous emergence holes to prevent egg-laying sites. <p>New houses usually infested by use of infested lumber. Adults may also come from firewood.</p> <p>Alternative controls for powder post beetles: small items, such as picture frames can be heated at 120 to 140 F for six hours to kill existing life stages. Freezing (0 F) infested wood for 72 hours will also kill all life stages. If all evidence indicates the infestation is localized, wood could be replaced. Watch for new holes in adjacent areas. Decrease moisture in wood through ventilation and moisture barriers. Central heat and air may reduce wood moisture so there is insufficient moisture to support large infestations in living areas. Wood kept below 14 percent moisture would be unsuitable to Anobiid powderpost beetle reinfestation or development. Professionals have moisture meters.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>RATS PB1624</p> <p>EPA has changed allowances regarding use of rodenticide baits in the urban environment. Check for latest updates before using baits.</p>	<p>Norway rat: 12-18 inches, tail shorter than head and body, body heavy and thick, ears small</p> <p>Roof rat: 12-17 inches, tail longer than head and body, body light and slender, ears larger.</p> <p>Young rat : 6-7 inches, feet large, head large</p> <p>House mouse: 6-7 inches feet small, head small</p> <p>Droppings:</p> <p>Roof rat: pointed, about ½ inch</p> <p>Norway rat: blunt, about ¾ inch</p> <p>House mouse: pointed, about 1/8 inch</p>	<p>When rats are plentiful or where unsanitary conditions exist with shelter, poisoned baits are the best control method. Often community- wide control needed. Poison baits are available as ready to use, premixed baits. They come in many forms: parafinized blocks for outdoor use and high humidity areas; treated meal; seeds; or parafinized pellets in bulk or in "place packs" for indoor use. Water baits are sold as packets of concentrate that are mixed with water. They are administered with a chick waterer and are useful in areas where rodent food is abundant. Poison baits should be placed where they are inaccessible to children and pets. Where rodent runs are exposed and in most outdoor situations, tamper proof bait boxes should be used and anchored. Vitamin K is the antidote for anti-coagulants.</p> <p>Second-Generation Anticoagulant Products for Professional Applicators must contain at least 16 pounds of bait. Bait stations are required for all outdoor, above-ground placements of second-generation anticoagulants. Bait stations are required indoors if exposure to children, pets, or non-target animals is possible. Distribution to and sales in "consumer" stores including grocery stores, drug stores, hardware stores, club stores will be prohibited. All outdoor above ground use must be in a bait station and be applied within 50 feet of buildings. (http://www.epa.gov/oppsrrd1/reregistration/rodenticides/finalriskdecision.htm)</p>	<p>Exclusion practices needed. Rats can fit through an opening ½ inch in diameter. Locate entrance into structure and exclude. Use materials such as galvanized, stainless or other non-rusting metal such as 24- gauge sheet metal or 19-gauge hardware cloth with 1/4 inch or smaller opening; brick, concrete block, tile or glass; steel wool or copper mesh with expandable foam; and others. Remove debris such as piles of waste lumber or trash, used feed sacks, abandoned large appliances and wood piles from next to structure. Store pet foods and seed in rodent proof glass or metal containers. Place snap traps, multiple catch traps and glue boards along paths traveled by rats. Of the snap traps, the expanded trigger trap is the most versatile since it can be baited. Rats are bait shy. Leave baits in place for at least a week before moving. Place trap 90 degrees to rodent pathway with trigger part of trap against the run. Rodents use edges of walls, studs and pipes as guidelines. Snap traps can be baited with: raisins or grapes for roof rats; sardines packed in oil for Norway rats; bacon squares; or small wads of cotton. Often area-wide effort needed.</p>
<p>SILVERFISH AND FIREBRATS SP341-O</p>	<p>Grayish, wingless, rapid-moving insects with 3 long tails. Feed on starchy materials such as bookbinding, wallpaper, cardboard, etc.</p>	<p>Niban Fine Granular Bait Maxforce Granular Insect Bait, Maxforce Fine Granule Insect Bait Tempo Ultra WP Cy-Kick Demand CS PT565 XLO PT Tri-Die Silica & Pyrethrum Dust (+Pressurized) Suspend SC DeltaDust Drione Talstar P</p>	<p>Treat crack and crevice where silverfish and firebrats may dwell. Attics many times source of infestation.</p>
<p>SKUNKS PB1624</p>	<p>These animals many times live in the ground around or under homes.</p>	<p>Bac-Azap biological odor control or others can be applied to eliminate odors.</p>	<p>Trap and remove skunks from property. Seal the foundation to prevent entry under building.</p>
<p>SNAILS AND SLUGS</p>	<p>Long, grayish, shiny, soft-bodied creatures. Will attack various plants. Leave slime trails on walks and walls.</p>	<p>Snail and slug killer baits containing metaldehyde.</p>	<p>Remove boards and plastic or plant debris and dry damp areas adjacent to foundation.</p>
<p>SNAKES PB1624</p>	<p>Snakes of various kinds, den around or invade homes and other buildings.</p>	<p>Place a pile of cool, damp rags in building where snake was last seen. Snake will be attracted and can be removed. Large glue boards can trap snakes.</p>	<p>Mouse-proof building. Mow lawns and field to control grass, weeds and brush. Remove boards, flat rocks, trash piles and other debris.</p>
<p>SOWBUGS OR PILLBUGS</p>	<p>Grayish, hard-shelled, many-legged creatures appear on walks and patios. Roll up in ball when disturbed. Occasional invaders.</p>	<p>Chemical control usually not necessary for this pest. If needed, apply to infested areas outdoors around perimeter of structure. This may stop any invasion into the house.</p> <p>Talstar P DeltaDust Suspend SC Tempo SC Ultra</p>	<p>Remove leaf piles, grass clippings, old boards, wood piles and other debris from around foundation. Leave a 12-18 inch plant /mulch free zone next to foundation base. Use exclusion practices: caulk cracks around foundation and screen vents in foundation. Drain and dry area around house.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TERMITES, SUBTERRANEAN PB 1344</p>	<p>Termites invade and eat wood and other cellulose material, causing extensive damage in structural parts of a building. Their presence may not be discovered until they swarm, years after infesting a structure.</p> <p>Workers light-colored, soft-bodied insects 1/8 to 3/16 inch long, soldier with a darkened head capsule.</p> <p>Swarms are black, brown or tan with wings. Wings are easily broken off after the mating flight and may be found by windows. Swarms are easily distinguished from winged ants by termite's straight antennae, broadly attached thorax to waist and four nearly equal wings. Subterranean termites with 2 darkened wing veins traveling the entire length of the fore wing.</p> <p>Inspect for signs of termite infestation: irregular earthen tubes constructed across walls, floors and foundation.</p> <p>Hammer or probe timbers with a sharp instrument. Damaged wood will be soft, channeled, unsound and may possibly reveal the termite infestation itself.</p> <p>Use a moisture meter. Active termites will increase moisture reading relative to uninfested areas.</p> <p>Termites commonly enter homes around doors, wooden steps and porches and unexcavated portions of structures. The easiest access points are where wood is in direct contact with the soil.</p> <p>(Description continued on next page)</p>	<p>Listed below are products available to professionals (Manufacturer):</p> <p>+ F = may also be foamed</p> <p>Soil treatment: Do not apply near (within 100 ft.) any body of water, cistern, or well.</p> <p>Nonrepellents (Newer a.i.s.) Termites do not detect these insecticides and walk over the treated soil. Termiticide may be transferred back to colony.</p> <p>Chlorantraniliprole (no signal word) Altriset 0.05% (DuPont) chlorfenapyr Phantom 0.125, 0.25% (BASF) + F fipronil Termidor HE 0.125% (BASF) Applications (2 gal/10 lin.ft., up to 2 ft deep, trench only 2 inch deep X 4 inch wide, 18 inch drill holes) differ from other liquid treatments, see Label for details. Termidor SC, 0.06%, 0.09%, 0.125% (BASF) +F Termidor 80WG, 0.06%, 0.09%, 0.125% (BASF)+F imidacloprid Premise 75, 0.05, 0.1% (Bayer)+F Premise 2 (small jobs and foaming) (Bayer)</p> <p>Pyrethroids (Older a.i.) In general, this groups tends to be repellent, thus treatments must be applied to create a continuous barrier. Many generic pyrethroids are now on the market.</p> <p>bifenthrin TalstarP 0.06, 0.12% (FMC) + F permethrin Dragnet SFR 0.5, 1, 2% (FMC) +F Prelude 0.5, 1, 2% (Amvac) +F</p> <p>Combination product Transport 0.11% bifenthrin and acetamiprid (FMC)</p> <p>A more complete list of termiticides is available at http://www.flaes.org/pdf/TermiticidesRegisteredInFlorida.pdf, but not all FL termiticides listed may be registered in Tennessee.</p> <p>(Wood treatment continued on next page)</p>	<p>Follow correct construction practices. This is the best protection against a termite infestation:</p> <ol style="list-style-type: none"> (1) Remove all wood materials from around and under the house; (2) Remove all form boards and construction stakes; (3) Construct a termite-proof foundation; (4) Have at least 30 in. of clearance under buildings; (5) Have proper ventilation and light under all parts of the building; (6) Use a moisture barrier in crawl spaces; (7) Drain water away from building; (8) Have no wood in contact with the ground, or treat those timbers that require ground contact with approved preservatives/borates (9) Make periodic inspection of buildings. <p>Find a reputable professional to treat.</p> <p>Collect some swarmer for identification and vacuum rest. Leave mud tubes in place until professional pest control applicator arrives.</p> <p>Newer Exterior Perimeter/Localized Interior treatment (EP/LIT) supplement to Termidor Labels. Ensure localized interior treatment is made. Label and more information at pestcontrolfacts.com. Premise also has an Exterior Perimeter/Interior Spot Treatment, see http://www.backedbybayer.com/system/product/product_label_pdf/65/Premise-75-432-1332-110322AV1-SRL.pdf for specifics.</p> <p>Tennessee Department of Agriculture now prefers the use of a disclosure form if less than a complete treatment is applied. Applications following the Exterior Perimeter/Interior treatments of Termidor or Premise are now considered a full treatment.</p> <p>New NPMA 33 Wood-destroying Insect Infestation report required for most real estate transactions. Sample form and directions are found at www.npmapestworld.org</p> <p>Control measures continued on the next page)</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TERMITES, SUBTERRANEAN Cont'd</p>	<p>Some termiticide labels have listed variable rates depending on soil type.</p> <p>To ensure your home is treated as directed by the label, ask the professional to provide you with a copy of the label.</p> <p>Suggested volumes listed below are usually for the lowest rate.</p> <p>In general, horizontal barriers (under slab) should receive 1 gallon of diluted termiticide per 10 square feet or 1.5 gallons if coarse fill.</p> <p>Vertical barriers (along both sides of foundation wall, around plumbing, piers and conduits) should receive 4 gallons of dilution per 10 linear feet per foot of depth (into a trench 6 inches wide) to the top of the footing not to exceed 4 ft.</p> <p>Voids in hollow masonry foundation walls should be treated at a rate of 2 gallons per 10 linear feet so the dilution will reach the top of the footing.</p> <p>Occasionally, moisture damaged wood in roofs can support an aerial infestation. No mud tubes will reach to ground. Attic inspection is important, too.</p>	<p>Wood treatment: treat galleries and wall voids; spray or brush on until wet, but NOT to runoff. Used to supplement a soil treatment. Most termiticides can be used in this manner.</p> <p>disodium octaborate tetrahydrate can be applied to wood as a pretreatment barrier or as a second barrier (see labels for sites and more details). Paint, spray, foam, inject or brush.</p> <p>Bora-Care (extensive label, other sites in addition to wood can be treated)</p> <p>Jecta (injection, for wood in contact with the ground) and others.</p> <p>Termidor Dry (0.5% fipronil) – To supplement other termite treatments. For termite galleries and shelter tubes apply 0.1 – 1g per injection point. For applications into carton and nests apply 0.3 - 3g per injection point. For applications into voids apply 0.2 - 2 g per injection point.</p> <p>Baits: Termites feed on bait and spread bait to colony to eliminate or suppress it. See PB1344 for advantages and disadvantages of baits.</p> <p>Sentricon® Colony Elimination System (DowAgrosciences LLC): Bait (Recruit IV and Recruit IV AG) contains a chitin synthesis inhibitor, noviflumuron. Bait Recruit HD allows once-a-year monitoring.</p> <p>Hex-Pro™ Termite Baiting System (DowAgrosciences LLC): Bait (Shatter) contains a chitin synthesis inhibitor, hexaflumuron.</p> <p>Exterra Termite Interception and Baiting System (Ensystem, Inc.) contains a chitin synthesis inhibitor, diflubenzuron.</p> <p>Advance™ Termite Baiting System (Whitmire-MicroGen) contains a chitin synthesis inhibitor, diflubenzuron.</p>	<p>Effective control measures for a soil treatment should include:</p> <ol style="list-style-type: none"> 1) Inspect basement and underside of house thoroughly to determine the area and extent of infestation. 2) Inspect attic for termite tubes and damage to joists, rafters, flooring and stored materials. 3) Disrupt and block all termite tubes (unless baiting) 4) Ditch the entire foundation inside and out and treat the soil replaced in the trenches with chemicals. 5) Repair all foundation and basement floor and wall breaks with rich concrete. 6) Break all wood-soil contacts, treat such areas with chemicals. 7) Treat infested timbers and replace those which are badly infested. 8) Treat hollow spaces in the foundation - concrete blocks, piers, chimney bases, spaces behind brick veneer. Drill and treat inside of porch foundations, under patios, under concrete slabs and the surface of ground under porches and similar dead places. 9) Provide ventilation and drainage beneath house and porches. 10) Remove all scrap wood from beneath house.
<p>TERMITES, DRYWOOD,</p>	<p>Swarmer with 3 darkened wing veins along the first third of the forewing.</p> <p>Soldier head capsule either rectangular with teeth on inner margin of left mandible or plug-shaped. All soldiers with pronotum wider than head capsule.</p> <p>Workers feed in and across the grain and leave six-sided fecal pellets piled below gallery openings. No mud present in galleries. Do not require connection to soil.</p>	<p>Spot or Localized Treatment Timbor (dust form) Termidor SC Termidor Foam Termidor Dry Others</p> <p>Whole Structure or Chamber Fumigation Sulfuryl Fluoride (requires structural fumigation license) Vikane Zythor Heat</p>	<p>Drywood termites may be controlled with a spot or localized treatment if the colony is small and accessible. If they are widely dispersed in structural lumbers, then a whole-house fumigation may be needed. Small infested items can be fumigated in a chamber.</p> <p>Although drywood termite distribution maps do not include Tennessee, several established populations are known from the Nashville area. Drywood termite infestations are most often introduced in furniture from Gulf coast states, California or other subtropical areas. Additional information can be found at</p> <p>http://www.floridatermitehelp.org/pdf/p00106_alternatedrywoodmethods_0910.pdf ; https://insects.tamu.edu/extension/publications/epubs/e-366.cfm;</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TICKS PB 726</p>	<p>brown or grey, oval to round, hard-shelled, 6 to 8-legged creatures which invade homes, yards and get on pets and people.</p>	<p>Insecticide applications are most effective when directed into areas where ticks and their animal hosts are likely to frequent. Pay particular attention to borders and fences between wooded or brushy areas and the lawn, around ornamental plantings, beside foot paths, house and dog house. Allow surface to dry before people or pets have access.</p> <p>Tempo SC Ultra Tempo Ultra WP Suspend DeltaGard G Talstar P Draagnet SFR Termiticide/Insecticide Astro and other synthetic pyrethroid insecticides.</p> <p><u>Indoors for brown dog tick:</u> Demand CS Talstar P Cy-Kick other pyrethroids</p> <p><u>Repellents:</u> Apply deet to skin; and Permanone 0.5% spray to shoes, cuffs and socks heed directions about drying before wearing.</p> <p><u>Dogs:</u> fipronil (Front Line) spot-ons available from veterinarians</p> <p>collars containing amitraz (Preventic) or flumethrin and imidacloprid (Seresto) (don't use around small children or dogs that may chew collar)</p> <p>BioSpot and other permethrin containing spot-ons</p> <p>See pesticide recommendations at http://www.ent.uga.edu/pmh/ under <i>Homeowner</i>, and <i>Animals</i>, for a thorough listing of veterinarian supplied on-pet products</p>	<p>Nonchemical methods for reducing tick problems include mowing the lawn and controlling weeds.</p> <p>This has three advantages - it lowers the moisture in the grass microclimate and allows sunlight to penetrate, which tends to cause ticks to dry out; it discourages rodents (which may serve as hosts) from nesting; and lastly, because there is less plant matter, less pesticide may be needed if a treatment is necessary. Also, removing debris, weeds or clutter from around the house discourages rodents from nesting.</p> <p>Repair entry points into the house to discourage possible tick hosts from entering. Cracks and crevices, both indoors and out, can be sealed to reduce hiding places for ticks. Inspect and clean pets and their bedding frequently. If bedding is infested, it can be cleaned or destroyed.</p> <p>In the home, ticks stay around baseboards and walls.</p> <p>Use insecticides in cracks and crevice in the home for brown dog tick.</p>
<p>WASPS, HORNETS, YELLOW-JACKETS SP290-A, SP341-M</p>	<p>Many types build paper and mud nests around homes, in ground or in shrubs.</p>	<p><u>Dusts:</u> Tempo 1D DeltaDust Drione Apicide</p> <p><u>Sprays:</u> Bee and wasp killer aerosols Tempo Ultra WP Tempo SC Ultra Eco PCO Jet</p> <p>Victor Yellow Jacket Trap</p>	<p>Wait until dark when wasps return to nest and are slow due to cooler temperatures. Apply insecticides to nest opening and seal nest opening if possible.</p> <p>Remove mud nests in winter to destroy overwintering forms.</p> <p>Traps can used to reduce foraging yellow jacket populations. Place away from areas people congregate.</p>

For Pest Management Professionals

Trade Name	Chemical Name or Use	Website
Abate Tire Treatment, Pellets	temephos	Clarke Mosquito Control http://www.clarke.com/
Advance Termite Bait System	diflubenzuron	BASF http://pestcontrol.basf.us/products/product-index.html
Alpine WSG	dinotefuran	BASF http://pestcontrol.basf.us/products/product-index.html
Altosid Briquets (B), Liquid Larvicide, Extended Residual Briquets (XR-B), Pellets (P), Extended Residual Granular (XR-G), Single Brood Granular (SBG) Pro G	(s) methoprene	Wellmark http://www.centralmosquitocontrol.com/
Anvil 10 + 10	sumithrin and PBO	Clarke Mosquito Control http://www.clarke.com/
Apicide	carbaryl	Mystic Chemical Company http://www.mysticchemical.com/images/labels/Apicide_Label.pdf
Aquabac xt, 200	Bacillus thuringiensis subspecies israelensis toxin	Becker Microbial Products http://beckermicrobialproductsinc.com/#!/products
Archer	pyriproxyfen	Syngenta http://www.syngentaprofessionalproducts.com/
Armor-Guard	disodium octaborate tetrahydrate	NovaGuard http://www.novaguard.com
Astro	permethrin	FMC http://pestsolutions.fmc.com/
Bac-Azap	enzyme-producing bacteria and others	Nisus http://www.nisuscorp.com/
Bedlam Plus	d-phenothrin. MGK-264, imidacloprid	McLaughlin, Gormley, King Co http://www.mgkpro.com
Biomist 3 + 15 ULV, 1.5 + 7.5 ULV, 30 + 30 ULV	permethrin and PBO	Clarke Mosquito Control http://www.clarke.com/
Bora-Care	disodium octaborate tetrahydrate	Nisus http://www.nisuscorp.com/
CimeXa	amorphous silica gel	Rockwell Labs http://www.rockwelllabs.com/
Cirkil	Neem oil	Terramera, Inc. http://cirkil.com/products , http://cirkil.com/rag
Cynoff EC	cypermethrin	FMC http://pestsolutions.fmc.com
CB-80	pyrethrin, PBO	FMC http://pestsolutions.fmc.com
DeltaDust	deltamethrin	Bayer http://www.backedbybayer.com/pest-management
DeltaGard G	deltamethrin	Bayer http://www.backedbybayer.com/pest-management
Demand CS	lambda cyhalothrin	Syngenta http://www.syngentaprofessionalproducts.com/
Demon WP	cypermethrin	Syngenta http://www.syngentaprofessionalproducts.com/
Dragnet SFR	permethrin	FMC http://pestsolutions.fmc.com/
Drione	silica gel and pyrethrins	Bayer http://www.backedbybayer.com/pest-management
DuPont Advion Ant Bait Arena	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html

Trade Name	Chemical Name or Use	Website
DuPont Advion Ant Gel	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Advion Cockroach Bait	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Advion Insect Granule	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Altriset Termiticide	chlorantraniliprole	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Arilon	indoxacarb	DuPont Professional Products http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
EcoPCO® D•X Dust Insecticide	2-phenethyl propionate & pyrethrins	http://www.envincio.com/products
EcoPCO Jet-	Eugenol, 2-phenethyl propionate	http://www.envincio.com/products
EcoRaider Bed Bug Killer	natural geraniol, cedar extract, sodium lauryl sulfate and natural derivative surfactant	EcoRaider http://ecoraiderusa.com/bedbugs.aspx
EndZone Insecticide Sticker	Acetamiprid	FMC http://pestsolutions.fmc.com/
Exterra Termite Interception & Baiting System	diflubenzuron	Ensystem http://www.ensystem.com/
Gentrol Aerosol, IGR concentrate	hydroprene	Wellmark- Zoecon http://www.zoecon.com/msdlabels.php
Gentrol Point Source	hydroprene	Wellmark- Zoecon http://www.zoecon.com/msdlabels.php
Golden Malrin Fly Bait	methomyl	Wellmark- Zoecon http://www.zoecon.com/msdlabels.php
Gourmet Ant Bait Gel Gourmet Ant Bait Liquid	disodium octaborate tetrahydrate	Innovative Pest Control Products http://www.antcafe.com/index.html
Intice Thiquid Ant Bait	1% borax	Rockwell labs http://www.rockwelllabs.com/
Jecta	10% disodium octaborate tetrahydrate	Nisus http://www.nisuscorp.com/
Kicker	pyrethrin and PBO	Bayer http://www.backedbybayer.com/pest-management
Kontrol 30-30	permethrin and PBO	Univar http://www.masterline.com/pdfs/Kontrol30-30Label122105.pdf
Mavrik Perimeter	tau-fluvalinate	Wellmark- Zoecon http://www.zoecon.com/msdlabels.php
Mattress Safe	bed encasement	http://www.mattresssafe.com/
Maxforce Ant Killer Bait Gel	fipronil	Bayer http://www.backedbybayer.com/pest-management
Maxforce Roach Killer Small Bait Stations	hydramethylnon	Bayer http://www.backedbybayer.com/pest-management
Maxforce Carpenter Ant Bait Gel	0.001% fipronil	Bayer http://www.backedbybayer.com/pest-management
Maxforce FC Magnum Roach Killer Bait Gel	0.05% fipronil	Bayer http://www.backedbybayer.com/pest-management
Maxforce FC Roach Bait Stations	fipronil	Bayer http://www.backedbybayer.com/pest-management
Maxforce FC Select Roach Killer Bait Gel	fipronil	Bayer http://www.backedbybayer.com/pest-management

Trade Name	Chemical Name or Use	Website
Maxforce Roach Killer Bait Gel	hydramethylnon	Bayer http://www.backedbybayer.com/pest-management
Maxforce Quantum Ant Bait	0.03% imidacloprid	Bayer http://www.backedbybayer.com/pest-management
MAXFORCE@COMPLETE Brand Granular insect Bait	1% hydramethylnon	Bayer http://www.backedbybayer.com/pest-management
Maxforce Granular Bait	hydramethylnon	Bayer http://www.backedbybayer.com/pest-management
Maxforce Granular Fly Bait	imidacloprid	Bayer http://www.backedbybayer.com/pest-management
Mosquitomist One	chlorpyrifos	Clarke Mosquito Control http://www.clarkemosquito.com
MotherEarth® D Pest Control Dust	diatomaceous earth	BASF http://pestcontrol.basf.us/products/product-index.htmlf
Mother Earth Granular Scatter Bait	boric acid	BASF http://pestcontrol.basf.us/products/product-index.htmlf
Natular 2E,G,G30, XRT and T30	spinosad	Clarke http://www.clarke.com/index.php
Niban FG = Niban Fine Granular Bait	orthoboric acid	Nisus http://www.nisuscorp.com/
Niban G = Niban Granular Bait	orthoboric acid	Nisus http://www.nisuscorp.com/
NiBor-D	disodium octaborate tetrahydrate	Nisus http://www.nisuscorp.com/
Nuvan Prostrip	dichlorvos	AMVAC http://www.amvac-chemical.com/Products/tabid/102/Default.aspx
Optigard Ant Gel Bait	thiamethoxam	Syngenta http://www.syngentaprofessionalproducts.com/
Pharorid - discontinued	methoprene	
Phantom,PT Phantom@II pressurized insecticide	chlorfenapyr	BASF http://pestcontrol.basf.us/products/product-index.html/
Precor Plus 2000 Premise Spray	permethrin, methoprene, phenothrin, etc.	Wellmark - Zoecon http://www.zoecon.com/msdlabels.php
Precor IGR Concentrate	methoprene	Wellmark - Zoecon http://www.zoecon.com/msdlabels.php
Premise 75, 2	imidacloprid	Bayer http://www.backedbybayer.com/pest-management/
Prelude	permethrin	AMVAC http://www.amvac-chemical.com/ContactUs/ProdDet/tabid/103/Default.aspx?pid=193
Prescription Treatment Advance 375A Select Granular Ant Bait	0.011% abamectin	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment 388B Advance Ant Bait Gel	5.4% sodium tetraborate decahydrate (borax)	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Advance Cockroach Gel Bait Reservoir	0.5% dinotefuran	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Alpine Dust	dinotefuran and diatomaceous earth	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Avert Dry Flowable Cockroach Bait Form 1	abamectin	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Cy-Kick, Cy-Kick CS	cyfluthrin	BASF http://pestcontrol.basf.us/products/product-index.html

Trade Name	Chemical Name or Use	Website
Prescription Treatment® Microcare® 3% CS Controlled Release Pyrethrins	pyrethrin, PBO + Other	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Tri-Die Dust, PT Tri-Die Pressurized	silica, pyrethrin, PBO	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment Ultracide	nylar (pyriproxyfen), pyrethrins, permethrin, etc.	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment® 221L Residual Insecticide	lambda-cyhalothrin	BASF http://pestcontrol.basf.us/products/product-index.html
Prescription Treatment® 565 Plus XLO Formula 2	pyrethrin, PBO + others	BASF http://pestcontrol.basf.us/products/product-index.html
Protect-A-Bed with BugLock™ 3 sided zipper system and ALLERZIP™ seal (bed encasement for bed bugs)	bed encasement	Protect-A-Bed http://www.protectabed.com http://www.protect-a-bed.com/
Recruit IV, Recruit IV AG, Recruit HD	noviflumuron	DowAgroSciences http://www.cdms.net/LabelsMsds/LMDefault.aspx?pd=7444
Scourge - Restricted Use Pesticide	resmethrin + PBO	Bayer http://www.backedbybayer.com
Sentricon Colony Elimination System	directions for use	DowAgroSciences http://www.dowagro.com/products/label/index.htm
Shatter	hexaflumuron	DowAgroSciences http://www.cdms.net/LabelsMsds/LMDefault.aspx?manuf=11&t=1,2,3,4
Steri-Fab	d-phenothrin isoproponal & others	Noble Pine Products Company http://www.sterifab.com/home.html
Suspend SC	deltamethrin	Bayer http://www.backedbybayer.com/pest-management/
Suspend Polyzone	deltamethrin	Bayer http://www.bayerprocentral.com/
Talstar P, PL	bifenthrin	FMC http://pestsolutions.fmc.com/
Tandem	thiomethoxam and lambda cyhalothrin	Syngenta http://www.syngentaprofessionalproducts.com/
Teknar HP-D, G	<i>Bacillus thuringiensis</i> subspecies <i>israelensis</i> toxin	Valent BioScience http://publichealth.valentbiosciences.com/products
Tempo Ultra WP	β-cyfluthrin	Bayer http://www.backedbybayer.com/pest-management
Tempo SC Ultra	β-cyfluthrin	Bayer http://www.backedbybayer.com/pest-management
Temprid SC, RTS	imidacloprid, β-cyfluthrin	Bayer http://www.backedbybayer.com/pest-management
Termidor Dry, for drywoods or to supplement other subterranean termite treatments	fipronil	BASF http://pestcontrol.basf.us/products/product-index.html
Termidor SC, 80 WG, HE, Foam	fipronil	BASF http://pestcontrol.basf.us/products/product-index.html
Terro-PCO Liquid Ant Bait	5.4% borax or sodium tetraborate decahydrate	Nisus http://www.nisuscorp.com/
Tim-bor	disodium octaborate tetrahydrate	Nisus http://www.nisuscorp.com/
Trade Name	Chemical Name or Use	Website

Transport GHP Insecticide	bifenthrin, acetamiprid	FMC http://pestsolutions.fmc.com/
Transport Termiticide Insecticide	bifenthrin, acetamiprid	FMC http://pestsolutions.fmc.com/
VectoBac GS, G, 12AS	<i>Bacillus thuringiensis</i> subspecies <i>israelensis</i> toxin	Valent BioScience http://publichealth.valentbiosciences.com/products
VectoLex CG, WSP, WDG	<i>Bacillus sphaericus</i>	Valent BioScience http://publichealth.valentbiosciences.com/products
Vendetta	abamectin B1	MGK http://www.mgk.com/professionalpestcontrol/
Vikane	sulfuryl fluoride	Douglas Products http://www.cdms.net/ldat/ld0KQ010.pdf
Zenprox EC	etofenprox and PBO	Zoecon http://www.zoecon.com/msdlabels.php
Zenprox Aerosol	etofenprox, tetramethrin, pyrethrins and PBO	Zoecon http://www.zoecon.com/msdlabels.php
Zyrox Fly Granular Bait	cyantraniliprole	Syngenta http://www.syngentapmp.com/
Zythor	sulfuryl fluoride	Ensystem http://www.ensystem.com/pdf/labels/Zythor%203-9.pdf

PRECAUTIONARY STATEMENT

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label.

DISCLAIMER STATEMENT

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label takes precedence over the recommendations found in this publication. Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), The University of Tennessee Institute of Agriculture and the University of Tennessee Extension assume no liability resulting from the use of these recommendations.