

Research Article

Methods for suspension culture, protoplast extraction, and transformation of high-biomass yielding perennial grass *Arundo donax*

Gaia Pigna^{1,2}, Taniya Dhillon¹, Elizabeth M. Dlugosz¹, Joshua S. Yuan⁴, Connor Gorman⁴, Piero Morandini^{2,3}, Scott C. Lenaghan^{5,6} and C. Neal Stewart, Jr.¹

¹ Department of Plant Sciences, University of Tennessee, Knoxville, USA

² Department of Biosciences, University of Milan, Milano, Italy

³ National Research Council, Institute of Biophysics, Milano, Italy

⁴ Department of Plant Pathology and Microbiology, Texas A&M University, College Station, TX, USA

⁵ Department of Food Science, University of Tennessee, Knoxville, USA

⁶ Department of Mechanical, Aerospace, and Biomedical Engineering, University of Tennessee, Knoxville, USA

Arundo donax L. is a promising biofuel feedstock in the Mediterranean region. Despite considerable interest in its genetic improvement, *Arundo* tissue culture and transformation remains arduous. The authors developed methodologies for cell- and tissue culture and genetic engineering in *Arundo*. A media screen was conducted, and a suspension culture was established using callus induced from stem axillary bud explants. DBAP medium, containing 9 μM 2,4-D and 4.4 μM BAP, was found to be the most effective medium among those tested for inducing cell suspension cultures, which resulted in a five-fold increase in tissue mass over 14 days. In contrast, CIM medium containing 13 μM 2,4-D, resulted in just a 1.4-fold increase in mass over the same period. Optimized suspension cultures were superior to previously-described solidified medium-based callus culture methods for tissue mass increase. Suspension cultures proved to be very effective for subsequent protoplast isolation. Protoplast electroporation resulted in a $3.3 \pm 1.5\%$ transformation efficiency. A dual fluorescent reporter gene vector enabled the direct comparison of the CAMV 35S promoter with the switchgrass *ubi2* promoter in single cells of *Arundo*. The switchgrass *ubi2* promoter resulted in noticeably higher reporter gene expression compared with that conferred by the 35S promoter in *Arundo*.

Received	15 AUG 2016
Revised	13 OCT 2016
Accepted	19 OCT 2016
Accepted article online	20 OCT 2016

Keywords: *Arundo* · Electroporation · Protoplast · Suspension Culture · Transformation

1 Introduction

In recent years there has been significant interest in harvesting renewable fuels and value-added chemicals from plant feedstocks, especially from perennial grasses such

as *Arundo donax* L. [1, 2, 3]. Some reasons why perennial grasses are desirable as next-generation bioenergy feedstock are: (i) their low soil management requirements and growth on marginal land; (ii) their ability to prevent soil erosion and increase soil organic carbon; (iii) few natural pests; and (iv) the capacity to reliably produce biomass [4]. In the USA, switchgrass (*Panicum virgatum*) is a widely adapted biomass feedstock whereas in Europe, *Arundo* may fulfill an important role as a perennial grass feedstock [5, 6]. Compared with switchgrass, *Arundo* has a much higher photosynthetic capacity, despite being a C3 plant [7, 8], leading to the production of significant biomass [9]. In addition, *Arundo* is salt tolerant, and has

Correspondence: C Neal Stewart, Jr., Department of Plant Sciences, University of Tennessee, 2431 Joe Johnson Dr., Knoxville, TN 37996, USA
E-mail: nealstewart@utk.edu

Abbreviations: BAP, benzylaminopurine; CIM, callus induction media; 2,4-D, 2,4-dichlorophenoxyacetic acid; GFP, green fluorescent protein; PEG, polyethylene glycol; PCV, packed cell volume

been used for phytoremediation of soils contaminated with heavy metals [10, 11]. Further, the cell wall composition of *Arundo* [12, 13], along with thermochemical conversion into biofuel [14], has been characterized. The lignin present in the stems and leaves has a HGS-type structure, but is primarily composed of G subunits. Delignified stems have a highly heterogeneous population of hemicelluloses in terms of sugar composition, molecular weight, and structure [13].

Despite the potential of *Arundo* as a bioenergy feedstock in Southern Europe [1], there are very few publications on tissue culture and transformation of this species. At present, *Agrobacterium*-mediated transformation of *Arundo* has not been demonstrated and limited success has been achieved using biolistics to transiently transform *Arundo* callus cultures [15]. A transient transformation protocol could provide crucial insight into gene expression [16], promoter screening [17], protein-protein interactions [18], transcriptional regulatory networks [19], and protein subcellular localization [20], as well as providing a necessary first step for the development of a stable transformation protocol for crop improvement [21]. As such, the goal of this work was to develop an in vitro culture system for *Arundo* to rapidly generate axenic protoplasts for subsequent transformation and analysis.

2 Materials and methods

2.1 Induction and maintenance of *Arundo* callus

Callus was induced using the protocol reported by Takahashi et al. [22] with some modification. Briefly, stem explants of *A. donax* were cut into approximately 30 cm long segments and surface sterilized with 70% v/v ethanol for 30 s and then transferred to a 4% v/v sodium hypochlorite solution, with one drop of Tween 20 (Sigma Aldrich, St. Louis, MO, USA), for 10 min prior to washing three times in sterilized distilled water. The cane internodes were then excised and placed on a basal Murashige and Skoog (MS) medium [23] supplemented with 0.05% Plant Preservative Mixture (PPM) (Plant Cell Technology, Washington, DC, USA) and 0.3% w/v Gelzan (Sigma Aldrich, St. Louis, MO, USA). The excised internodes were placed in a growth chamber at 25°C under standard cool white fluorescent tubes with a flux rate of 35 $\mu\text{mol s}^{-1} \text{m}^{-2}$ and a 16-h photoperiod. Excised internodes were transferred onto CIM medium, MS medium supplemented with 3% sucrose (Fisher Scientific, Fair Lawn, NJ, USA), 9 μM 2,4-dichlorophenoxyacetic acid (2,4-D), 0.7 g/L casein hydrolysate, 0.5 g/L L-glutamine, 1 g/L L-proline, and 0.4% Gelzan. These were placed in a growth chamber in the dark at 25°C for a month to induce calli. Once induced, the calli were maintained on the same medium, with 13 μM 2,4-D [22].

2.2 Establishment and maintenance of *Arundo* suspension cultures

Cell suspension cultures were initiated from 10 month old embryogenic callus by adding 500 mg of calli to a 125 mL Erlenmeyer flask containing 25 mL of MS or B5 medium supplemented with 3% sucrose and various hormones. In order to test the effect of different media on initiation of suspension cultures, the hormone type and concentration were varied, along with the basal media as follows. DBAP medium was composed of MS salts, B5 vitamins, 9 μM 2,4-D and 4.4 μM 6-benzylaminopurine (BAP) (Sigma Aldrich, St. Louis, MO, USA) [24]. MSDP media was composed of MS salts and vitamins supplemented with 9 μM 2,4-D and 100 nM phyto-sulfokine- α (PSK- α) (PolyPeptide Group, Torrance CA, USA) [25]. B5G medium was composed of B5 salts and vitamins [26] supplemented with 0.7 g/L glutamine (Sigma Aldrich, St. Louis, MO, USA) [27]. After transfer of the calli to flasks, cultures were incubated in the dark at 28°C for 14 days on a gyratory shaker at 80 rpm. Cultures were then filtered through a 100 μm Fisherbrand™ Cell Strainer (Fisher Scientific, Fair Lawn, NJ, USA), brought up to a volume of 25 mL with fresh medium, and transferred to a 125 mL flask. Three flasks, initiated separately from three different calli, were used for each medium and represented three independent biological replicates. Five milliliters of fresh medium was added to the cultures every seven days for six to eight weeks to compensate liquid evaporation. At the end of eight weeks, cultures were considered to be established, as indicated by the presence of large cell clusters [28, 29] and the growth was assayed for another 21 days by measuring the packed cell volume (PCV) of the cultures [30]. Briefly, 5 mL of culture was transferred to a 15 mL conical centrifuge tube (Corning Incorporated) and allowed to settle for 10 min, at which point the PCV was measured. At the end of the experiments, the cultures were used to generate protoplasts for subsequent experiments.

2.3 Tissue culture compared with cell suspension culture

In order to determine the best strategy for rapidly attaining large amounts of *Arundo* tissue, we tested four culture conditions, with three independent replicates for each condition, all of which were initiated with 500 mg (fresh weight) of callus or cells. Treatment 1 (liquid DBAP to liquid DBAP) was initiated with 500 mg of an established suspension culture added to 25 mL of DBAP medium in an Erlenmeyer flask. Each replicate was initiated from a different flask of suspension cell culture. Treatment 2 (liquid DBAP to solid CIM) was initiated the same as Treatment 1, with the exception of the established suspension culture being added to solid CIM media. Treatment 3 (solid CIM to liquid DBAP) was initiated from callus grown on solid CIM media transferred directly into 25 mL of

DBAP medium in an Erlenmeyer flask. Finally, Treatment 4 (solid CIM to solid CIM) was initiated from callus grown on solid CIM media transferred to solid CIM media. To determine the tissue mass at 1, 3, 5, 7, 10, 14, 18, and 21 days, the suspension cultures were filtered through a 70 µm Fisherbrand™ Cell Strainer (Fisher Scientific, Fair Lawn, NJ, USA) and the contents weighed, accounting for the mass of the cell strainer, and then returned to the culture. The mass of the calli grown on solid media, at the same time points, was measured by transferring calli to a Petri dish and taking fresh weight.

2.4 Protoplast isolation from suspension cell culture

Protoplasts of *Arundo* were isolated using a previously established method for isolation of protoplasts from switchgrass suspension cultures [31] with slight modifications. In addition to using switchgrass as a control, tobacco (*Nicotiana tabacum* L.) cv. Bright Yellow 2 (BY-2) suspension culture, cultured as previously described, was used as a positive control to ensure functionality of the enzymes [31, 32, 33]. Briefly, a 10 mL PCV of *Arundo* suspension culture collected during the exponential growth phase was used. Cell walls were digested with 20 mL of a solution (0.6 M mannitol, 10 mM MES pH 5, 1 mM CaCl₂, 0.1% BSA, 5 mM 2-mercaptoethanol) containing food grade enzymes (Rohament CL 15820 ECU, Rohapect 10L 10080 ADJU, and Rohapect UF 0.078 ADJU) (AB Enzymes, Darmstadt, Germany) which was added to the cells and incubated at 37°C, for 2–3 h with gentle shaking. After incubation, the solution was filtered through a 40 µm Fisherbrand™ cell strainer to remove cellular debris and centrifuged at 150 g for 10 min. Five volumes of W5 wash solution (154 mM NaCl, 125 mM CaCl₂, 5 mM KCl, 2 mM MES, pH 5.7) [34, 35] was then added to the protoplasts. After resuspension, protoplasts were imaged using an Olympus IX 81 inverted microscope and counted using a hemocytometer. Protoplast diameter was measured using the ImageJ software package (US National Institutes of Health, Bethesda, Maryland, USA) calibrated with an ocular micrometer. The size distribution of the protoplasts was generated by grouping protoplasts in different size classes [36].

2.5 Vector construction

Vector pTD-PvUbi2-35S was derived from pANIC10A [37] after several modifications. First, a unique restriction site, *Mlu*I, was inserted between the switchgrass polyubiquitin 1+3 (*PvUbi1+3*) promoter and the orange fluorescent protein, *pporRFP*. Since *PvUbi2* is a stronger promoter compared to *PvUbi1+3* [38], the former was PCR amplified from pANIC10A and used to replace *PvUbi1+3* using *Psp*XI and *Mlu*I-HF restriction enzymes. To compare the strength of the constitutive cauliflower mosaic virus 35S promoter relative to *PvUbi2* in *A. donax*, a 35S-*mGFP5ER*-

nos cassette was PCR amplified from pBIN m-gfp5-ER [39] using the forward primer 5'-taatctgcttggtaccagattagccttccaat-3' and reverse primer 5'-ttacttggtggcgcgcctcccgatctagtaacataga-3' (restriction sites are underlined) and inserted in pANIC10A using the restriction enzymes *Bst*EII and *Asc*I. The primers used for PCR amplification were designed in SnapGene and synthesized by Integrated DNA Technologies (Coralville, IA, USA). PCR was performed using Platinum *Taq* DNA polymerase (ThermoFisher Scientific, Waltham, MA, USA). To improve the efficiency of protoplast transformation, the size of the original vector was reduced by deleting unneeded sequences including (i) a 3837 bp fragment encompassing 85% of the total length of ZmUbi1-Gateway-compatible *ccdB* cassette-OCS, using *Mfe*I, (ii) a 3405 bp sequence containing DNA required for *Agrobacterium*-mediated transformation, which included the right T-DNA border, the stability region, and 652 bp of the origin of replication, using *Fsp*I, and (iii) a 2977 bp sequence spanning *PvUbi2* promoter and hygromycin selectable marker, using *Pac*I and *Psp*XI. All PCR amplification and restriction digestion products were gel purified using the QIAquick gel extraction kit (QIAGEN, Hilden, Germany) and quantified on a NanoDrop-1000 spectrophotometer (ThermoFisher) prior to cloning. Following *Pac*I and *Psp*XI digestion, the vector was blunt-ended and 5' phosphorylated using the Quick Blunting Kit (New England Biolabs, Ipswich, MA, USA). All ligation reactions were carried out using T4 DNA Ligase (New England Biolabs). Ligated constructs were transformed into either "One Shot *ccdB* survival 2 T1R" chemically competent cells or "One Shot Mach1 T1 phage-resistant" chemically competent cells (ThermoFisher) as per the manufacturer's instructions and sequence confirmed. The final vector, pTD-PvUbi2-35S contains 8690 bp (Fig. 1) and features two independent fluorescent protein cassettes: the orange fluorescent protein *pporRFP* under the control of *PvUbi2* promoter and the green fluorescent protein *mGFP5-ER* under the control of CAMV 35S promoter.

2.6 Polyethylene glycol (PEG)-mediated transformation

PEG-mediated DNA transformation was performed as previously described [21, 35], with switchgrass protoplasts serving as a positive control, with slight modifications. Protoplasts of *Arundo* were resuspended in MMg (0.4 M mannitol, 25–150 mM MgCl₂, 4 mM MES, pH 5.7) at a concentration of 1 × 10⁶ protoplasts/mL. 10 µg of pANIC10A-GFPstuffer [38] plasmid DNA was then mixed with 200 µL of protoplasts (approximately 2 × 10⁵ protoplasts). A 0–40% PEG solution (0.6 M mannitol, 100 mM CaCl₂, 0–50% PEG 4000 (Sigma-Aldrich, St. Louis, Missouri, USA) was added to the protoplasts to a final PEG concentration of 0, 10 or 20%. After 20 min of incubation at room temperature (RT), protoplasts were washed twice




Figure 1. Map of pTD-PvUbi2-35S vector. Abbreviations: *PvUbi2*, *Panicum virgatum* polyubiquitin 2 promoter and intron; *pporRFP*, *Porites porites* red fluorescent protein coding region; *nosT*, *Agrobacterium tumefaciens* nopaline synthase terminator CaMV 35S, cauliflower mosaic virus 35S promoter; *mgfp5-ER*, modified green fluorescent protein with an endoplasmic reticulum targeting sequence; *ColE1*, origin of replication in *E. coli*; *bom*, origin of transfer in *E. coli*; *KanR*, kanamycin bacterial resistance marker; *LB*, T-DNA left border. Unique restriction sites used for vector construction are also indicated.

with 1 and 4 mL of W5 and collected by centrifugation at 100 $\times g$ for 6 min. Protoplasts were then resuspended in 1 mL of WI (0.6 M mannitol, 4 mM KCl, 4 mM MES, pH 5.7), transferred to a 12-well plate (Corning Incorporated, Corning, NY, USA) and incubated at 28°C in the dark for 20 h. As a positive control, switchgrass mesophyll protoplasts were transformed using PEG as previously described [31]. The transformation efficiency (%) was calculated by counting the number of protoplasts expressing *pporRFP* divided by the total number of protoplasts using a hemocytometer and multiplied by 100 as shown in the formula below:

$$\left(\frac{\# \text{ protoplasts expressing } pporRFP}{\text{total \# protoplasts}} \right) \times 100 =$$

% transformation efficiency

To ensure the detection of all positive protoplasts, samples were collected from individual wells, centrifuged at 100 $\times g$ prior to resuspension in a minimal volume (~100 μL) and counted on a hemocytometer. Three independent biological replicates were analyzed for each tested condition.

2.7 Protoplast electroporation-mediated transformation

As an alternative to PEG-mediated transformation, electroporation was performed as previously described for

tobacco BY-2 protoplasts with slight modifications [40]. 1 h after isolation, protoplasts were centrifuged at 100 $\times g$ for 2 min and the W5 wash solution was removed. Protoplasts were then washed twice with electroporation buffer (0.4 M sucrose, 2.4 g/L HEPES, 6 g/L KCl, 600 mg/L $\text{CaCl}_2 \cdot \text{H}_2\text{O}$, pH 7.2), at 80 $\times g$ for 10 min without deceleration. A 500 μL aliquot containing 5 to 6 $\times 10^5$ protoplasts in electroporation buffer was then placed into 0.4 cm gap electroporation cuvette (Bio-Rad Laboratories, Hercules, CA, USA). Forty micrograms of pTD-PvUbi2-35S DNA was then added to *Arundo* and tobacco BY-2 protoplasts and incubated at RT for 10–15 min, placing the cuvette horizontally. After incubation, electroporation of *Arundo* was conducted using a Bio-Rad Gene Pulser™ (Bio-Rad Laboratories, Hercules, CA, USA) at 130–300 V with a capacitance of 500 or 1000 μF . Tobacco BY-2 protoplasts were used as a positive control, at 300 V with a capacitance of 1000 μF . The time constant was recorded after each pulse to ensure the samples were uniform, and the pulse was delivered. After electroporation, samples were incubated for 40 min at RT in the cuvette, and then moved into a single well of a 12-well plate, followed by the addition of 2 mL of protoplast culture medium (4.3 g/L MS salts, 0.4 M sucrose, 2.5 mM MES hydrate, 5.4 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$, 3.1 mM NH_4NO_3 , pH 5.7). The remaining solution in the cuvette was mixed with 1 mL of protoplast culture medium and transferred into a second well. An additional 1 mL of protoplast culture medium was finally used to wash the cuvette and poured into the second well. The protoplast viability was monitored after electroporation using propidium iodide, working solution (1 mg/50 mL, Sigma–Aldrich, St. Louis, Missouri, USA) [31]. Similarly to PEG-mediated transformation, the transformation efficiency was measured at 14–16 h after transformation by counting the number of protoplasts expressing *mGFP5-ER* and *pporRFP* and dividing by the total protoplast number. For this method efficiency of transformation was estimated directly in the well by counting three random fields of view. The efficiency of each field was calculated separately as described previously. Using this strategy, an average of about 300 protoplasts were counted per well. Three independent biological replicates, each the result of independent protoplast isolations and transformation, were analyzed for all conditions tested.

2.8 Statistical analysis

A completely random experimental design was used for media screening, growth analysis, and transformation. All experiments included at least three independent biological replicates, and were analyzed using mixed model ANOVA (IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY, USA). Least significant differences (LSD) were used to determine significant differences among means when the ANOVA results were statistically significant ($P < 0.05$).

3 Results

3.1 Establishment of suspension cultures of *Arundo*

To test various media, the PCV of cultures were analyzed over the course of 21 days (Fig. 2A). Despite using solidified CIM medium for culturing the donor callus, liquid CIM resulted in the poorest growth among the media tested (Table 1). Over the first seven days, the PCV of the liquid CIM was below the limit of detection (50 μ L), and reached a measurable level only at day 10. Moreover, there was no significant growth observed for the CIM cultures from day 10 to day 21 ($p = 0.71$). Similar to CIM, suspension cultures initiated in MSDP medium showed no detectable growth over the first 10 days of culture, but reached a measurable level at day 14. From day 14 to the completion of the experiments at day 21, there was no significant growth ($p = 0.13$). Significant growth was observed in the B5G and DBAP media, with the cultures reaching their peak growth at 7–10 days and 10–14 days respectively (Table 1). In B5G and DBAP, the PCV was measurable after only one day in culture, as opposed to three and seven days for MSDP and CIM respectively. Between DBAP and B5G, there was no significant difference in growth over the first three days of culture ($p = 0.65$); however, the PCV of the B5G culture was significantly greater at day 7 compared to DBAP ($p = 0.024$). After 10 days in culture, the PCV of the suspension cultures in DBAP had increased 3.3-fold, while the B5G culture increased 2.8-fold compared with the initial amount. At this point, the DBAP culture surpassed the maximum PCV observed for the B5G culture. Furthermore, comparison of the PCV at day 10 (the maximum for both DBAP and B5G) across all of the media showed a significant difference between DBAP and B5G compared to the other two media tested (Table 1). After 10 days in culture, the PCV of the B5G culture decreased markedly. A similar trend was observed in the DBAP culture, but the decrease occurred after 14 days in culture. In both cases, the rapid cell growth in these cultures led to an observable decrease in the amount of liquid media in flasks. While we conclude that both DBAP and B5G were suitable for initiation and maintenance of *Arundo* cell suspension cultures, DBAP was chosen as the media for future experiments because of its higher PCV at 10 days of culture, indicating more biomass would be available sooner for follow-on experiments.

Prior to establishment of the cultures, the majority of the cells in DBAP were kidney-shaped, contained starch-rich plastids, and clearly-discernable nuclei (Fig. 2B). After establishment, the suspension cultures mainly consisted of elongated cells devoid of vacuoles (Fig. 2C). In addition, the established suspension cultures were a mix of individual embryogenic-like cells of various size classes (Fig. 2E). Once large (1–5 mm diameter) clusters were readily observable in the culture flasks, the cultures were considered suitable for subsequent experiments.




Figure 2. Suspension cell culture establishment and protoplast isolation. (A) One month-old callus grown on solid CIM medium. (B) Cells from seven day-old filtrated suspension cell culture in liquid DBAP medium. (C) Cells from 14 day-old established suspension cell culture in DBAP medium. (D) Callus from one month-old suspension cell culture regrown on CIM solidified medium. (E) Large cell clusters (1–5 mm) after one week of subculture of 5 mL PCV in 20 mL fresh DBAP medium in a 125 mL Erlenmeyer flask. (F) Experimental scheme for comparing tissue culture and suspension cell culture. All treatments were initiated from callus on solid CIM: (1) liquid DBAP to liquid DBAP; (2) liquid DBAP to solid CIM; (3) solid CIM to liquid DBAP; (4) solid CIM to solidified CIM medium. (G) Protoplasts isolated from a five day-old suspension cell culture. (H) Distribution of protoplast size in five samples of isolated protoplasts. Scale bar in (A) and (D) is 500 μ m, scale bar in (B), (C) and (G) is 50 μ m.

3.2 *Arundo* performs better in liquid culture than on solidified medium

A comparison of the mass of suspension cells and callus was also evaluated over the course of 21 days (Table 2, Fig. 2F). At 14 days of culture, there was no significant difference in the amount of tissue generated from *Arundo* grown on solid media (maintained on solid CIM), a freshly incubated liquid culture (callus moved from solid CIM to liquid DBAP medium), or an established liquid culture transferred to solid media (liquid DBAP to solid DBAP) (Fig. 2D). However, there was significantly more

Table 1. Media screen for the establishment of suspension cultures over a 21-day culture period. Cell growth was monitored for 21 days after filtration through a 100 µm cell strainer by measuring the packed cell volume (PCV). Three independent replicates were measured for each medium.

Media	Packed cell volume (µL) measured by day					
	1	3	7	10	14	21
DBAP	900 ± 10	100 ± 20	130 ± 30	280 ± 30	280 ± 30 (a) ^(a)	90 ± 10
CIM	N.D. ^(b)	N.D.	N.D.	60 ± 20	60 ± 10 (b)	60 ± 40
MSDP	N.D.	N.D.	N.D.	N.D.	60 ± 20 (b)	200 ± 100
B5G	80	90 ± 10	220 ± 30	220 ± 30	120 ± 30 (b)	90 ± 10

(a) Letter designations at 14 days indicates the results of Tukey's HSD analysis, with common letters among treatments representing no significant difference from three biological replicates.

(b) N.D. indicates that the packed cell volume was below the detection limit (50 µL).

Table 2. Comparison of various cell- and tissue culture regimes to generate tissue mass over a 21-day culture period. The starting material for each replicate was 500 mg with the treatment time course (day 0) beginning with the second medium in each treatment. Three independent replicates were measured for each treatment.

Treatments	Fresh weight (mg) per day							
	1	3	5	7	10	14	18	21
Liquid DBAP to liquid DBAP	500 ± 10	700 ± 70	730 ± 50	940 ± 70	1500 ± 130	2400 ± 380 (a) ^(a)	2300 ± 570	2000 ± 600
Solid CIM to solid CIM	510 ± 10	540 ± 30	580 ± 40	630 ± 10	730 ± 70	720 ± 20 (b)	720 ± 210	730 ± 200
Solid CIM to liquid DBAP	510 ± 10	720 ± 30	740 ± 150	780 ± 180	750 ± 230	740 ± 240 (b)	830 ± 290	660 ± 280
Liquid DBAP to solid CIM	500	660 ± 50	720 ± 40	820 ± 20	860 ± 90	940 ± 140 (b)	980 ± 70	970 ± 80

(a) Letter designations at 14 days indicates the results of Tukey's HSD analysis, with common letters between treatments representing no significant difference from three biological replicates.

tissue generated from the established suspension cultures that were transferred to fresh liquid DBAP. In fact, 2.5 times more tissue was generated by cultures maintained in liquid DBAP compared to all other treatments at 14 days (Table 2). In addition, the established suspension cultures reached a maximum of 4.7 times the starting tissue mass over 21 days, whereas no other treatment exceeded two times the cell mass over the same time period. Based on these results, the established suspension cultures were used for the generation of protoplasts and transformation.

3.3 Protoplast preparation and transformation.

Protoplasts of *Arundo* were isolated from four to seven day old suspension cell cultures (Fig. 2G), with an average of $3.8 \times 10^5 \pm 2.5 \times 10^5$ protoplasts/mL of PCV. Morphologically, the protoplasts ranged in size from 4.9 to 53.4 µm in diameter with an average size of 20.7 ± 12.3 µm. The majority of the protoplasts ranged from 4 to 20 µm in diameter (Fig. 2H). PEG- and electroporation transformation experiments were performed on protoplasts. Initial transformation experiments were conducted using the pANI-C10A-GFPstuffer [31], which expresses *pporRFP* driven by the switchgrass ubiquitin promoter, *PvUbi1+3* [38]. All

attempts to transform *Arundo* protoplasts with PEG (varying DNA quantity, $MgCl_2$ concentration, PEG% solution) failed, despite the success with the switchgrass positive controls. However, electroporation at 130 V and 1000 µF attained a transformation efficiency of $3.3 \pm 1.5\%$. When the voltage was increased to 300 V, with the capacitance remaining the same, there was a decrease in protoplast viability from $67.7 \pm 3.0\%$ to $11.9 \pm 4.5\%$ and no transformation was detected. When the capacitance was decreased to 500 µF, at 130 and 300 V, the efficiency of transformation was reduced to 1.0 ± 0.2 and $0.3 \pm 0.4\%$, respectively. In these treatments, the viability remained high, $82 \pm 1.5\%$ at 130 V and $80 \pm 2.4\%$ at 300 V, despite the lower transformation efficiency. Not surprisingly, the switchgrass *PvUbi2* [38] promoter yielded higher marker gene expression than the *35S* promoter in *Arundo* protoplasts using the pTD-*PvUbi2*-*35S* binary vector (Fig. 3A-D). As expected, tobacco BY-2 protoplasts, used as a control, showed high expression of the mGFP5-ER fluorescent marker driven by the *35S* promoter, but only weak expression of the *pporRFP* fluorescent marker driven by the *PvUbi2* promoter (Fig. 3E-H).




Figure 3. Transient expression of GFP and pporRFP fluorescent proteins in *Arundo* and tobacco BY-2 protoplasts transformed with pTD-*PvUbi2-35S* by electroporation. (A, B, C, D) *Arundo* protoplast. (E, F, G, H) tobacco BY-2 protoplast. Visualization of transformation using a TdTomato filter set: 545/30 nm excitation and 605/50 nm band pass emission (A, B, E, F) and GFP filter set: 470/30 nm excitation and 525/50 nm band pass emission (C, D, G, H). (B, D, F, H) represent the merge of fluorescent and bright field images. Exposure time was 300 ms and the scale bar is 50 μ m in all pictures.

4 Discussion

Previous work on tissue culture of *Arundo donax* used solid CIM medium for callus induction and maintenance [22]. Unfortunately, *Arundo* callus grown on this medium is relatively slow-growing, making it difficult and time-consuming to generate enough tissue for subsequent transformation. Of particular importance, *Arundo* callus growth on CIM appears to be suboptimal compared with other relevant plant species. For example, *Sorghum* callus grow 1.7 times faster than *Arundo* on solidified CIM [41]. For these reasons, a suspension culture system for *Arundo* was pursued as a mean to increase the amount of tissue produced. The results from the *Arundo* suspension cultures established in DBAP demonstrate that cultivation in liquid medium produces far more tissue than solid culture with CIM. Similar results have been found in other species, such as cotton (*Gossypium hirsutum* L.) and broccoli (*Brassica oleracea*) [42, 43]. The difference in *Arundo* growth in liquid culture may be the result of greater nutrient availability, a more gradual pH change, or lower production of toxic compounds [42].

Despite the significant increase in tissue mass for the suspension cultures of *Arundo*, the amount obtained was still relatively low when compared to suspension cultures of other biofuel crops. Suspension cultures of sugarcane (*Saccharum officinarum*) have been shown to increase by six times the starting inoculum in only 10 days [44]. Similarly, switchgrass suspension cultures have been shown to increase 8.6 times over 14 days of culture [24]. In both of these systems, the increase in fresh weight is about two times greater than with *Arundo*. These data suggest that further optimization of the culture media may be necessary for *Arundo*, as for other bioenergy crops researchers have placed extensive effort in this direction [45, 46, 47]. Also, the ability of plant species to be manipulated in vitro is strongly genotype-dependent [48, 49], therefore more effort is needed to screen different accessions and identify genotypes more amenable to in vitro cultivation.

One unique characteristic of *Arundo* compared with other bioenergy crops is that it does not sexually reproduce; therefore there is no seed production [50, 51]. As such, *Arundo* is not amenable to improvement through standard breeding practices. The lack of seed production, however, is a highly advantageous trait when considering bioconfinement and restriction of gene flow, especially transgene flow [52]. In either case, genetic improvement of *Arundo* requires a reliable method for transformation to introduce novel genes, or inhibit/overexpress native ones [53, 54]. An effective transformation method is also essential for screening of gene expression, validation of gene silencing or genome-editing vectors, and general cell and molecular assays. Considering that only there have been few RNA sequencing studies [55, 56] and no whole genome information is presently available for *Arundo*, it is essential that screening be conducted at early stages, prior to developing transgenic plants. Currently, transformation of *Arundo* callus has only been achieved in one study, in which particle bombardment was used to introduce exogenous DNA into callus [15]. While effective, particle bombardment requires specialized equipment and is costly with regards to consumables. Further complicating transient-, and to a greater extent, stable transformation, callus growth is very slow, leading to a long lag time prior to when screening can be performed. As an alternative, PEG-mediated transformation and electroporation of *Arundo* protoplasts was evaluated in our project. High molecular weight PEG solutions have proven to be toxic to protoplasts isolated from some plant species [57]. In addition, the presence of Ca^{2+} or Mg^{2+} and the ratio of cations/PEG are critical factors for optimizing protoplast transformation [58, 59]. While we did vary these parameters, we never achieved PEG-mediated transformation of *Arundo* protoplasts. However, while PEG-mediated transformation failed, electroporation was successful in generating transient expression of the fluorescent reporter genes. While, similar to particle bombardment, electroporation requires specialized equipment, its throughput can be markedly increased using plate-based assays. In addi-

tion, screening after electroporation can be conducted in less than 24 h, providing a rapid assessment of gene expression. To demonstrate the functionality of the *Arundo* protoplast transformation system, the expression of fluorescent reporters driven by monocot (*PvUbi2*) and dicot (*35S*) promoters were evaluated using the pTD-*PvUbi2-35S* vector developed in this work. The results demonstrated the utility of screening promoters/gene expression in a rapid protoplast system: *35S* promoter imbued low expression in *Arundo* protoplasts. This finding was in contrast to previous work using particle bombardment to transform *Arundo* callus [15]. However, protoplast systems are much more sensitive for measuring fluorescent protein expression compared with callus-based systems [29]. The identification of *PvUbi2* as a strong promoter for *Arundo* has implications in the development of future vectors aimed at the generation of stably-transformed *Arundo*. For instance, a strong promoter should be chosen to drive the plant resistance cassette in order to impose a strong selection and eliminate the majority of false positives/escapes [60, 61]. Assays such as the one conducted in this work may help to overcome current limitations in the generation of transgenic *Arundo*, by providing data on promoter activity prior to regeneration [62].

In conclusion, in our project, a cell suspension culture system using DBAP medium was established for *Arundo*, which yielded up to six-fold tissue-mass increase relative to callus culture systems. In addition, we developed a low-cost protoplast isolation system from suspension cultures that appear to have utility for assays for gene expression, and promoter screening. They could be useful for genome editing and stable transformation. Finally, the transient transformation of *Arundo* protoplasts using electroporation was achieved and used to screen a representative monocot and dicot promoter. These experiments indicated that *PvUbi2* was a strongly-active promoter in *Arundo*, whereas *35S* was a weak promoter, thus informing the design of future vectors for *Arundo* biotechnology.

This work was supported by ARPA-E award number DE-AR0000313. The authors thank Robert N. Trigiano for the use of the Bio-Rad Gene Pulser™.


The authors declare no financial or commercial conflict of interest.

5 References

- [1] Lewandowski, I., Scurlock, J. M., Lindvall, E., Christou, M., The development and current status of perennial rhizomatous grasses as energy crops in the US and Europe. *Biomass Bioenergy* 2003, 25, 335–361.
- [2] Corno, L., Pilu, R., Tran, K., Tambone, F. et al., Sugars production for green chemistry from 2nd generation crop (*Arundo donax* L.): A full field approach. *ChemistrySelect* 2016, 1, 2617–2623.
- [3] Corno, L., Pilu, R., Cantaluppi, E., Adani, F., Giant cane (*Arundo donax* L.) for biogas production: The effect of two ensilage methods on biomass characteristics and biogas potential. *Biomass Bioenergy* 2016, 93, 131–136.
- [4] Angelini, L. G., Ceccarini, L., Di Nasso, N. N., Bonari, E., Comparison of *Arundo donax* L. and *Miscanthus x giganteus* in a long-term field experiment in Central Italy: Analysis of productive characteristics and energy balance. *Biomass Bioenergy* 2009, 33, 635–643.
- [5] Mantineo, M., D'agosta, G. M., Copani, V., Patanè, C., Cosentino, S. L., Biomass yield and energy balance of three perennial crops for energy use in the semi-arid Mediterranean environment. *Field Crops Res.* 2009, 114, 204–213.
- [6] Scordia, D., Testa, G., Cosentino, S. L., Perennial grasses as lignocellulosic feedstock for second-generation bioethanol production in Mediterranean environment. *Ital. J. Agron.* 2014, 9, 84–92.
- [7] Rossa, B., Tüffers, A. V., Naidoo, G., Willert, V. D., *Arundo donax* L. (Poaceae) – a C3 species with unusually high photosynthetic capacity. *Bot. Acta* 1998, 111, 216–221.
- [8] Ceotto, E., Di Candilo, M., Castelli, F., Badeck, F.-W. et al., Comparing solar radiation interception and use efficiency for the energy crops giant reed (*Arundo donax* L.) and sweet sorghum (*Sorghum bicolor* L. Moench). *Field Crops Res.* 2013, 149, 159–166.
- [9] Beale, C. V., Long, S. P., Can perennial C4 grasses attain high efficiencies of radiant energy conversion in cool climates? *Plant Cell Environ.* 1995, 18, 641–650.
- [10] Papazoglou, E. G., Karantounias, G. A., Vemmos, S. N., Bouranis, D. L., Photosynthesis and growth responses of giant reed (*Arundo donax* L.) to the heavy metals Cd and Ni. *Environ. Int.* 2005, 31, 243–249.
- [11] Mirza, N., Mahmood, Q., Pervez, A., Ahmad, R. et al., Phytoremediation potential of *Arundo donax* in arsenic-contaminated synthetic wastewater. *Bioresour. Technol.* 2010, 101, 5815–5819.
- [12] You, T. T., Mao, J. Z., Yuan, T. Q., Wen, J. L., Xu, F., Structural elucidation of the lignins from stems and foliage of *Arundo donax* Linn. *J. Agric. Food Chem.* 2013, 61, 5361–5370.
- [13] Peng, F., Bian, J., Peng, P., Xiao, H., Ren, J. L. et al., Separation and characterization of acetyl and non-acetyl hemicelluloses of *Arundo donax* by ammonium sulfate precipitation. *J. Agric. Food Chem.* 2012, 60, 4039–4047.
- [14] Basso, M. C., Cerrella, E. G., Buonomo, E. L., Bonelli, P. R., Cukierman, A. L., Thermochemical conversion of *Arundo donax* into useful solid products. *Energy Sources* 2005, 27, 1429–1438.
- [15] Dhir, S., Knowles, K., Pagan, C. L., Mann, J., Optimization and transformation of *Arundo donax* L. using particle bombardment. *Afr. J. Biotechnol.* 2010, 9, 6460–6469.
- [16] Chen, J., Yi, Q., Song, Q., Gu, Y. et al., A highly efficient maize nucleus protoplast system for transient gene expression and studying programmed cell death-related processes. *Plant Cell Rep.* 2015, 34, 1239–1251.
- [17] Nakashima, K., Jan, A., Todaka, D., Maruyama, K. et al., Comparative functional analysis of six drought-responsive promoters in transgenic rice. *Planta* 2014, 239, 47–60.
- [18] Fujikawa, Y., Nakanishi, T., Kawakami, H., Yamasaki, K. et al., Split luciferase complementation assay to detect regulated protein-protein interactions in rice protoplasts in a large-scale format. *Rice* 2014, 7, 1.
- [19] Lin, Y. C., Li, W., Sun, Y. H., Kumari, S. et al., SND1 transcription factor-directed quantitative functional hierarchical genetic regulatory network in wood formation in *Populus trichocarpa*. *Plant Cell* 2013, 25, pp.4324–4341.

- [20] Nieves-Cordones, M., Chavanieu, A., Jeanguenin, L., Alcon, C. et al., Distinct amino acids in the C-linker domain of the *Arabidopsis* K⁺-channel KAT2 determine its subcellular localization and activity at the plasma membrane. *Plant Physiol.* 2014, *164*, 1415–1429.
- [21] Masani, M. Y. A., Noll, G. A., Parveez, G. K. A., Sambanthamurthi, R., Prüfer, D., Efficient transformation of oil palm protoplasts by PEG-mediated transfection and DNA microinjection. *PLoS One* 2014, *9*, e96831.
- [22] Takahashi, W., Takamizo, T., Kobayashi, M., Ebina, M., Plant regeneration from calli in giant reed (*Arundo donax* L.). *Grassland Sci.* 2010, *56*, 224–229.
- [23] Murashige, T., Skoog, F., A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant.* 1962, *15*, 473–497.
- [24] Gupta, S. D., Conger, B. V., Somatic embryogenesis and plant regeneration from suspension cultures of switchgrass. *Crop Sci.* 1999, *39*, 243–247.
- [25] Eun, C. H., Ko, S. M., Matsubayashi, Y., Sakagami, Y., Kamada, H., Phytosulfokine- α requires auxin to stimulate carrot non-embryogenic cell proliferation. *Plant Physiol. Biochem.* 2003, *41*, 447–452.
- [26] Gamborg, O. L., Miller, R., Ojima, K., Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell Res.* 1968, *50*, 151–158.
- [27] Finer, J. J., Nagasawa, A., Development of an embryogenic suspension culture of soybean (*Glycine max* Merrill.). *Plant Cell Tissue Org. Cult.* 1988, *15*, 125–136.
- [28] Savoure, A., Magyar, Z., Pierre, M., Brown, S. et al., Activation of the cell cycle machinery and the isoflavonoid biosynthesis pathway by active *Rhizobium meliloti* Nod signal molecules in *Medicago* microcallus suspensions. *EMBO J.* 1994, *13*, 1093.
- [29] Cole, M., Jacobs, B., Soubigou-Tacconat, L., Balzergue, S. et al., Live imaging of DORNROSCHEN and DORNROSCHEN-LIKE promoter activity reveals dynamic changes in cell identity at the microcallus surface of *Arabidopsis* embryonic suspensions. *Plant Cell Rep.* 2013, *32*, pp.45–59.
- [30] Mustafa, N. R., de Winter, W., van Iren, F., Verpoorte, R., Initiation, growth and cryopreservation of plant cell suspension cultures. *Nat. Protoc.* 2011, *6*, 715–742.
- [31] Burris, K. P., Dlugosz, E. M., Collins, A. G., Stewart Jr, C. N., Lenaghan, S. C., Development of a rapid, low-cost protoplast transfection system for switchgrass (*Panicum virgatum* L.). *Plant Cell Rep.* 2016, *35*, 693–704.
- [32] Buntru, M., Vogel, S., Spiegel, H., Schillberg, S., Tobacco BY-2 cell-free lysate: An alternative and highly-productive plant-based *in vitro* translation system. *BMC Biotechnol.* 2014, *14*, 1.
- [33] Buntru, M., Vogel, S., Stoff, K., Spiegel, H., Schillberg, S., A versatile coupled cell-free transcription–translation system based on tobacco BY-2 cell lysates. *Biotechnol. Bioeng.* 2015, *112*, 867–878.
- [34] Sheen, J., Signal transduction in maize and *Arabidopsis* mesophyll protoplasts. *Plant Physiol.* 2001, *127*, 1466–1475.
- [35] Yoo, S. D., Cho, Y. H., Sheen, J., *Arabidopsis* mesophyll protoplasts: A versatile cell system for transient gene expression analysis. *Nature Protoc.* 2007, *2*, 1565–1572.
- [36] Kim, G. H., Klotchkova, T. A., Kang, Y. M., Life without a cell membrane: Regeneration of protoplasts from disintegrated cells of the marine green alga *Bryopsis plumosa*. *J. Cell Sci.* 2001, *114*, 2009–2014.
- [37] Mann, D. G., LaFayette, P. R., Abercrombie, L. L., King, Z. R. et al., Gateway-compatible vectors for high-throughput gene functional analysis in switchgrass (*Panicum virgatum* L.) and other monocot species. *Plant Biotechnol. J.* 2012, *10*, 226–236.
- [38] Mann, D. G., King, Z. R., Liu, W., Joyce, B. L. et al., Switchgrass (*Panicum virgatum* L.) polyubiquitin gene (PvUbi1 and PvUbi2) promoters for use in plant transformation. *BMC Biotechnol.* 2011, *11*, 74.
- [39] Haseloff, J., Siemering, K. R., Prasher, D. C., Hodge, S., Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic *Arabidopsis* plants brightly. *Proc. Natl. Acad. Sci. USA* 1997, *94*, 2122–2127.
- [40] Miao, Y., Jiang, L., Transient expression of fluorescent fusion proteins in protoplasts of suspension cultured cells. *Nat. Protoc.* 2007, *2*, 2348–2353.
- [41] Hagio, T., Blowers, A. D., Earle, E. D., Stable transformation of sorghum cell cultures after bombardment with DNA-coated microprojectiles. *Plant Cell Rep.* 1991, *10*, 260–264.
- [42] Gawel, N. J., Robacker, C. D., Somatic embryogenesis in two *Gossypium hirsutum* genotypes on semi-solid versus liquid proliferation media. *Plant Cell Tissue Org. Cult.* 1990, *23*, 201–204.
- [43] Yang, J. L., Seong, E. S., Kim, M. J., Ghimire, B. K. et al., Direct somatic embryogenesis from pericycle cells of broccoli (*Brassica oleracea* L. var. italica) root explants. *Plant Cell Tissue Org. Cult.* 2010, *100*, 49–58.
- [44] Ho, W. J., Vasil, I. K., Somatic embryogenesis in sugarcane (*Saccharum officinarum* L.): Growth and plant regeneration from embryogenic cell suspension cultures. *Ann. Bot.* 1983, *51*, 719–726.
- [45] Perera, D., Barnes, D. J., Baldwin, B. S., Reichert, N. A., Direct and indirect *in vitro* regeneration of *Miscanthus* \times *giganteus* cultivar Freedom: Effects of explant type and medium on regeneration efficiency. *In Vitro Cell. Dev. Biol. Plant* 2015, *51*, 294–302.
- [46] Liu, Y. R., Cen, H. F., Yan, J. P., Zhang, Y. W., Zhang, W. J., Inside out: High-efficiency plant regeneration and *Agrobacterium*-mediated transformation of upland and lowland switchgrass cultivars. *Plant Cell Rep.* 2015, *34*, 1099–1108.
- [47] Birch, R. G., Sugarcane biotechnology: Axenic culture, gene transfer, and transgene expression, in: Moore, P. H., Botha, F. C., (Ed.), *Sugarcane: Physiology, Biochemistry, and Functional Biology*, John Wiley & Sons Ltd. 2013, pp. 645–681.
- [48] Tomita, K. I., Hiura, S., Tamagake, H., Evaluation of the potential for somatic embryogenesis in sugar beet (*Beta vulgaris* L.) breeding lines and improvement of regeneration efficiency. *Plant Biotechnol.* 2013, *30*, 479–487.
- [49] Gurel, E., Gurel, S., Lemaux, P. G., Biotechnology applications for sugar beet. *Crit. Rev. Plant Sci.* 2008, *27*, 108–140.
- [50] Hardion, L., Verlaque, R., Rosato, M., Rossello, J. A., Vila, B., Impact of polyploidy on fertility variation of Mediterranean *Arundo* L. (Poaceae). *C. R. Biol.* 2015, *338*, pp. 298–306.
- [51] Hardion, L., Verlaque, R., Baumel, A., Juin, M., Vila, B., Revised systematics of Mediterranean *Arundo* (Poaceae) based on AFLP fingerprints and morphology. *Taxon* 2012, *61*, 1217–1226.
- [52] Barney, J. N., DiTomaso, J. M., Nonnative species and bioenergy: Are we cultivating the next invader?. *Bioscience* 2008, *58*, 64–70.
- [53] Biswal, A. K., Hao, Z., Pattathil, S., Yang, X. et al., Downregulation of GAUT12 in *Populus deltoides* by RNA silencing results in reduced recalcitrance, increased growth and reduced xylan and pectin in a woody biofuel feedstock. *Biotechnol. Biofuels* 2015, *8*, p.1.
- [54] Jung, J. H., Altpeter, F., TALEN mediated targeted mutagenesis of the caffeic acid O-methyltransferase in highly polyploid sugarcane improves cell wall composition for production of bioethanol. *Plant Mol. Biol.* 2016, *92*, 131–142.
- [55] Fu, Y., Poli, M., Sablok, G., Wang, B. et al., Dissection of early transcriptional responses to water stress in *Arundo donax* L. by unigene-based RNA-seq. *Biotechnol. Biofuels* 2016, *9*, 54.
- [56] Barrero, R. A., Guerrero, F. D., Moolhuijzen, P., Goolsby, J. A. et al., 2015. Shoot transcriptome of the giant reed, *Arundo donax*. *Data Brief* 2015, *3*, 1–6.

- [57] Kao, K. N., Saleem, M., Improved fusion of mesophyll and cotyledon protoplasts with PEG and high pH-Ca²⁺ solutions. *J. Plant Physiol.* 1986, *122*, 217–225.
- [58] Lazzeri, P. A., Brettschneider, R., Lührs, R., Lörz, H., Stable transformation of barley via PEG-induced direct DNA uptake into protoplasts. *Theor. Appl. Genet.* 1991, *81*, 437–444.
- [59] Negrutiu, I., Shillito, R., Potrykus, I., Biasini, G., Sala, F., 1987. Hybrid genes in the analysis of transformation conditions. *Plant Mol. Biol.* 1987, *8*, 363–373.
- [60] Cao, J., Duan, X., McElroy, D., Wu, R., Regeneration of herbicide resistant transgenic rice plants following microprojectile-mediated transformation of suspension culture cells. *Plant Cell Rep.* 1992, *11*, 586–591.
- [61] Christou, P., Ford, T. L., Kofron, M., Production of transgenic rice (*Oryza sativa* L.) plants from agronomically important indica and japonica varieties via electric discharge particle acceleration of exogenous DNA into immature zygotic embryos. *Nat. Biotechnol.* 1991, *9*, 957–962.
- [62] Takahashi, W., Takamizo, T., *Molecular breeding of grasses by transgenic approaches for biofuel production*. INTECH Open Access Publisher 2012.



Cover illustration

This issue focuses on the ECAB conference and is edited by Professors Raquel Barros, Jochen Büchs, and Guilherme Ferreira. Transmission electron micrograph (circle) of metabolically engineered *E. coli* cells accumulating polyhydroxyalkanoates as white distinct granules. Poly(2-hydroxyisovalerate-co-lactate) structure out of the cell is schematically shown. The Image is provided by Jung Eun Yang, Je Woong Kim, Young Hoon Oh, So Young Choi, Hyuk Lee, A-Reum Park, Jihoon Shin, Si Jae Par, and Sang Yup Lee authors of "Biosynthesis of poly(2-hydroxyisovalerate-co-lactate) by metabolically engineered *Escherichia coli*". Cover image created by Jung Eun Yang and Sang Yup Lee (<http://dx.doi.org/10.1002/biot.201600420>).

Inside cover illustration

This issue focuses on the ECAB conference and is edited by Professors Raquel Barros, Jochen Büchs, and Guilherme Ferreira. Existing purification technologies for retroviral vectors result in very low yields of infective particles, often associated with the absence or loss of the envelope protein at the surface of viral particles. A peptide selective for amphotropic viral particles was discovered by panning a phage display peptide library, and employed to selectively capture and recover bound particles under gentle conditions. The Image is provided by Cláudia S. M. Fernandes, Inês Barbosa, Rute Castro, Ana Sofia Pina, Ana Sofia Coroadinha, Ana Barbas, and A. Cecília A. Roque authors of "Retroviral particles are effectively purified on an affinity matrix containing peptides selected by phage-display" (<http://dx.doi.org/10.1002/biot.201600025>).

Biotechnology Journal – list of articles published in the December 2016 issue.

Editorial

ECAB focus issue: Engineered catalysts, robust, cost-effective and integrated bioprocesses and high-throughput screening
Raquel Aires-Barros, Jochen Buechs, Guilherme Ferreira
<http://dx.doi.org/10.1002/biot.201600662>

Forum

Intellectual property rights, standards and data exchange in systems biology
Esther van Zimmeren, Berthold Rutz, Timo Minssen
<http://dx.doi.org/10.1002/biot.201600109>

Forum

Book review: An Introduction to Biomechanics Solids and Fluids, Analysis and Design
Ümit Keskin
<http://dx.doi.org/10.1002/biot.201600610>

Review

Extreme makeover: Engineering the activity of a thermostable alcohol dehydrogenase (AdhD) from *Pyrococcus furiosus*
Kusum Solanki, Walaa Abdallah and Scott Banta
<http://dx.doi.org/10.1002/biot.201600152>

Review

Miniaturization of aqueous two-phase extraction for biological applications: From micro-tubes to microchannels
Ruben R. G. Soares, Daniel F. C. Silva, Pedro Fernandes, Ana M. Azevedo, Virginia Chu, João P. Conde and M. Raquel Aires-Barros
<http://dx.doi.org/10.1002/biot.201600356>

Research Article

Retroviral particles are effectively purified on an affinity matrix containing peptides selected by phage-display
Cláudia S. M. Fernandes, Inês Barbosa, Rute Castro, Ana Sofia Pina, Ana Sofia Coroadinha, Ana Barbas and A. Cecília A. Roque
<http://dx.doi.org/10.1002/biot.201600025>

Research Article

Permeability of currently available microtiter plate sealing tapes fail to fulfil the requirements for aerobic microbial cultivation
Michaela Sieben, Heiner Giese, Jan-Hendrik Grosch, Kira Kauffmann and Jochen Büchs
<http://dx.doi.org/10.1002/biot.201600054>

Research Article

Mixing at the microscale: Power input in shaken microtiter plates
Astrid Dürauer, Stefanie Hobiger, Cornelia Walther and Alois Jungbauer
<http://dx.doi.org/10.1002/biot.201600027>

Review

The potential of nanoparticles in stem cell differentiation and further therapeutic applications
Ahmed Abdal Dayem, Hye Yeon Choi, Gwang-Mo Yang, Kyeongseok Kim, Subbroto Kumar Saha, Jin-Hoi Kim, and Ssang-Goo Cho
<http://dx.doi.org/10.1002/biot.201600453>

Mini-review

Integrated fabrication-conjugation methods for polymeric and hybrid microparticles for programmable drug delivery and biosensing applications
Sukwon Jung, Chang-Hyung Choi, Chang-Soo Lee and Hyunmin Yi
<http://dx.doi.org/10.1002/biot.201500298>

Research Article

Biosynthesis of poly(2-hydroxyisovalerate-co-lactate) by metabolically engineered *Escherichia coli*
Jung Eun Yang, Je Woong Kim, Young Hoon Oh, So Young Choi, Hyuk Lee, A-Reum Park, Jihoon Shin, Si Jae Park and Sang Yup Lee
<http://dx.doi.org/10.1002/biot.201600420>

Research Article

Recombinant flavin-dependent halogenases are functional in tobacco chloroplasts without co-expression of flavin reductase genes

Sabine Fräbel, Markus Krischke, Agata Staniek and Heribert Warzecha

<http://dx.doi.org/10.1002/biot.201600337>

Research Article

Engineering *Halomonas* spp. as a low-cost production host for production of bio-surfactant protein PhaP

Lu-Hong Lan, Han Zhao, Jin-Chun Chen and Guo-Qiang Chen

<http://dx.doi.org/10.1002/biot.201600459>

Research Article

Parallel online multi-wavelength (2D) fluorescence spectroscopy in each well of a continuously shaken microtiter plate

Tobias Ladner, Mario Beckers, Bernd Hitzmann and Jochen Büchs

<http://dx.doi.org/10.1002/biot.201600515>

Research Article

Microheterogeneity of therapeutic monoclonal antibodies is governed by changes in the surface charge of the protein

Beate Hintersteiner, Nico Lingg, Evelyne Janzek, Oliver Mutschlechner, Hans Loibner and Alois Jungbauer

<http://dx.doi.org/10.1002/biot.201600504>

Research Article

Scaling up a chemically-defined aggregate-based suspension culture system for neural commitment of human pluripotent stem cells

Cláudia C. Miranda, Tiago G. Fernandes, M. Margarida Diogo and Joaquim M. S. Cabral

<http://dx.doi.org/10.1002/biot.201600446>

Research Article

Screening through the PLICable promoter toolbox enhances protein production in *Escherichia coli*

Jianhua Yang, Anna Joëlle Ruff, Stefanie Nicole Hamer, Feng Cheng and Ulrich Schwaneberg

<http://dx.doi.org/10.1002/biot.201600270>

Research Article

Case study on human α 1-antitrypsin: Recombinant protein titers obtained by commercial ELISA kits are inaccurate

Henning Gram Hansen, Helene Faustrup Kildegaard, Gyun Min Lee and Stefan Kol

<http://dx.doi.org/10.1002/biot.201600409>

Research Article

Methods for suspension culture, protoplast extraction, and transformation of high-biomass yielding perennial grass *Arundo donax*

Gaia Pigna, Taniya Dhillon, Elizabeth M. Dlugosz, Joshua S. Yuan, Connor Gorman, Piero Morandini, Scott C. Lenaghan and C. Neal Stewart, Jr.

<http://dx.doi.org/10.1002/biot.201600486>

Biotech Method

Impedance technology reduces the enumeration time of *Brettanomyces* yeast during beer fermentation

Sanelle van Wyk and Filipa V. M. Silva

<http://dx.doi.org/10.1002/biot.201600497>

Biotech Method

Rotation-based technique for the rapid densification of tubular collagen gel scaffolds

Caroline Loy, Audrey Lainé and Diego Mantovani

<http://dx.doi.org/10.1002/biot.201600268>