

# Hybridization of downregulated-COMT transgenic switchgrass lines with field-selected switchgrass for improved biomass traits

Holly L. Baxter · Lisa W. Alexander · Mitra Mazarei · Ellen Haynes ·  
Geoffrey B. Turner · Robert W. Sykes · Stephen R. Decker · Mark F. Davis ·  
Richard A. Dixon · Zeng-Yu Wang · C. Neal Stewart Jr.

Received: 3 September 2015 / Accepted: 31 December 2015 / Published online: 21 January 2016  
© Springer Science+Business Media Dordrecht 2016

**Abstract** Transgenic switchgrass (*Panicum virgatum* L.) has been produced for improved cell walls for biofuels. For instance, downregulated caffeic acid 3-*O*-methyltransferase (COMT) switchgrass produced significantly more biomass and biofuel than the non-transgenic progenitor line. In the present study we sought to further improve biomass characteristics by crossing the downregulated COMT T<sub>1</sub> lines with high-yielding switchgrass accessions in two genetic backgrounds (‘Alamo’ and ‘Kanlow’). Crosses and T<sub>2</sub> progeny analyses were made under greenhouse conditions to assess maternal effects, plant morphology and yield, and cell wall traits. Female parent type influenced morphology, but had no effect on cell wall traits. T<sub>2</sub> hybrids produced with T<sub>1</sub> COMT-

downregulated switchgrass as the female parent were taller, produced more tillers, and produced 63 % more biomass compared with those produced using the field selected accession as the female parent. Transgene status (presence or absence of transgene) influenced both growth and cell wall traits. T<sub>2</sub> transgenic hybrids were 7 % shorter 80 days after sowing and produced 43 % less biomass than non-transgenic null-segregant hybrids. Cell wall-related differences included lower lignin content, reduced syringyl-to-guaiacyl (S/G) lignin monomer ratio, and a 12 % increase in total sugar release in the T<sub>2</sub> transgenic hybrids compared to non-transgenic null segregants. This is the first study to evaluate the feasibility of transferring the low-recalcitrance traits associated with a transgenic switchgrass line into high-yielding field varieties in an attempt to improve growth-related traits. Our

---

Holly L. Baxter and Lisa W. Alexander have equally contributed.

---

H. L. Baxter · L. W. Alexander · M. Mazarei ·  
E. Haynes · C. Neal Stewart Jr. (✉)  
Department of Plant Sciences, University of Tennessee,  
Knoxville, TN 37996, USA  
e-mail: nealstewart@utk.edu

H. L. Baxter · M. Mazarei · G. B. Turner ·  
R. W. Sykes · S. R. Decker · M. F. Davis ·  
R. A. Dixon · Z.-Y. Wang · C. Neal Stewart Jr.  
Oak Ridge National Laboratory, BioEnergy Science  
Center (BESC), Oak Ridge, TN 37831, USA

L. W. Alexander  
Otis L. Floyd Nursery Research Center, US National  
Arboretum, USDA-ARS, McMinnville, TN 37110, USA

G. B. Turner · R. W. Sykes · S. R. Decker · M. F. Davis  
National Renewable Energy Laboratory, Golden,  
CO 80401, USA

R. A. Dixon  
Department of Biological Sciences, University of North  
Texas, Denton, TX 76203, USA

Z.-Y. Wang  
Samuel Roberts Noble Foundation, Ardmore, OK 73401,  
USA

results provide insights into the possible improvement of switchgrass productivity via biotechnology paired with plant breeding.

**Keywords** COMT · Switchgrass · Lignocellulosic biofuel · Transgenic · Hybridization

## Introduction

Switchgrass (*Panicum virgatum*, L.), a warm-season C<sub>4</sub> perennial grass species native to North America, has significant potential as a renewable source of fuel because of its adaptation to a wide range of climates, high biomass potential on marginal land, and high nutrient- and water-use efficiency (Parrish and Fike 2005). In addition, switchgrass reduces soil erosion, improves water quality, and its deep root system enhances soil carbon sequestration and improves soil quality (McLaughlin and Walsh 1998; Schmer et al. 2008). As a highly self-incompatible species, switchgrass is considered an obligate outcrosser and relies on wind-mediated pollination for fertilization (Ge et al. 2011). Despite favorable economic and environmental traits, large-scale deployment of switchgrass and other lignocellulosic feedstocks as sustainable and cost-competitive sources of fuel will rely on improvements to feedstock quality, biomass productivity, and other desirable agronomic traits. In particular, recent breeding and biotechnology efforts with switchgrass have focused on improving biomass composition for bioconversion (Fu et al. 2011, 2012; Sarath et al. 2008; Shen et al. 2012) as well as enhancing plant architecture and/or biomass yield (Poovaiah et al. 2015; Vogel et al. 2013; Wuddineh et al. 2015a, b).

The efficiency of converting lignocellulosic biomass into biofuel depends largely on the chemical composition of plant cell walls, particularly the lignin content and composition (Chen and Dixon 2007). Lignin, a heterogeneous phenolic polymer comprised of *p*-hydroxyphenyl (H), guaiacyl (G) and syringyl (S) monomer units, interacts with cell wall polysaccharides to form a complex structure that is resistant to enzymatic breakdown (Boerjan et al. 2003). Several greenhouse and field experiments have shown that cell wall saccharification and biofuel yields of switchgrass and other biofuel feedstocks can be significantly improved by manipulating the lignin biosynthetic

pathway (Baxter et al. 2015; Poovaiah et al. 2014). One approach is to reduce the expression of a single gene in the lignin biosynthetic pathway, as demonstrated by switchgrass plants with downregulated caffeic acid *O*-methyltransferase (COMT), an enzyme primarily involved in S-lignin synthesis. Greenhouse and field experiments show that COMT-downregulated switchgrass lines have reduced lignin content and lower S/G lignin monomer ratios, higher sugar release by enzymatic hydrolysis, and up to 38 % higher ethanol yields compared with controls (Fu et al. 2011; Baxter et al. 2014).

We hypothesized that hybrid plants from crosses of altered-cell wall transgenic switchgrass lines with locally-adapted, high biomass accessions would result in high-biomass lines with improved bioconversion efficiency; yet, there are no published studies in this area. With this in mind, T<sub>1</sub>-generation COMT-downregulated switchgrass in an ‘Alamo’ genetic background was hybridized with ‘Alamo’ and ‘Kanlow’ accessions from local Tennessee fields based on superior growth performance, and the resulting T<sub>2</sub> hybrid progeny from these crosses were evaluated. The objectives of this study were to (1) evaluate relationships among seed weight, germination rate, and early morphology in T<sub>2</sub> hybrid families, (2) assess the influence of female parent type (T<sub>1</sub> COMT transgenic or field accession) on growth and cell wall traits among T<sub>2</sub> hybrid families, (3) compare plant growth and cell wall traits between transgenic and null segregant (non-transgenic) T<sub>2</sub> hybrids, and (4) examine the relationships among early morphology, reproductive stage morphology, dry weight, and cell wall traits (lignin content, S/G ratios, and total sugar release) in T<sub>2</sub> hybrids.

## Materials and methods

### Generation and maintenance of hybrid families

Twenty-six accessions from ‘Alamo’ and ‘Kanlow’ (lowland switchgrass cultivars) were selected for high biomass, plant height, and vigor from research or production fields in eastern Tennessee in 2010 and propagated in the greenhouse. Transgenic low-recalcitrance lines were vegetatively propagated from two T<sub>1</sub>-generation transgenic events (COMT2 and COMT3) that were produced in a previous study by

RNAi-mediated gene silencing of the caffeic acid *O*-methyltransferase (COMT) gene in ‘Alamo’ seed-derived switchgrass (Fu et al. 2011). In that previous study, two independent  $T_0$  events were crossed to the same ‘Alamo’ genotype to produce the  $T_1$ -generation COMT2 and COMT3 events. Thus,  $T_1$  COMT2 and  $T_1$  COMT3 are considered ‘Alamo’ half-siblings.  $T_1$  COMT transgenic parents and field-selected clones used for the crosses were maintained in 12 L pots, watered as needed and fertilized every 2 weeks with  $100 \text{ mg L}^{-1}$  of Peter’s 20-20-20 N-P-K fertilizer. Full-sibling and half-sibling  $T_2$ -generation hybrid switchgrass families were produced from crosses between transgenic low-recalcitrant lines and field accessions of high-yielding switchgrass.

Flowers were pollinated as they became receptive. Panicles were considered receptive when the distal-most floret opened. Receptive panicles were emasculated and bagged using DelNet pollination bags ( $508 \text{ mm} \times 457.2 \text{ mm}$ ; DelStar Technologies, Inc., Middletown, Delaware). Transgenic parents were crossed to wild type field selection parents using either open pollination to generate the half-sibling hybrid families, or mutual pollination to generate full-sibling hybrid families. Mutual pollinations were performed in a greenhouse (16 h photoperiod;  $27^\circ \text{C}$  day temperature and  $21^\circ \text{C}$  night temperature). For mutual pollinations, one or two receptive panicles from each parent plant were placed in a single bag, secured with a twist-tie, labeled, and staked to prevent damage. Crosses made using the mutual pollination method produced full-sibling families under greenhouse or walk-in growth chamber conditions.

Bags were removed for seed harvest after 90 days and dehisced seed in pollination bags was discarded. Panicles were separated and the seeds were threshed by hand on a rubber board. Seeds were counted, weighed, and treated with powdered ethyl mercaptan (CAPTAN<sup>®</sup>), and stored in airtight plastic bags at  $4^\circ \text{C}$  for a minimum of 30 days. In total, 9641 seeds were collected from 62 controlled crosses and 94,688 open-pollinated seeds were collected from 35 individual plants (26 wild type field selections and 9  $T_1$  COMT transgenics) for a total of 62 full-sibling and 35 half-sibling  $T_2$  hybrid families. Of these  $T_2$  hybrid families, 30 (20 full-sibling and 10 half sibling) were used for the current experiments.


Eighteen seeds per family were sown in Fafard 3B potting mix in  $7.6 \times 7.6 \times 5.7 \text{ cm}$  containers and maintained in the greenhouse. Germination frequency was determined as the total number of seedlings above the soil line 30 days after sowing (DAS) divided by the total number of seed sown per family. An average of 14 seedlings per family germinated. Seedlings were watered as needed. After the appearance of two true leaves, the seedlings were fertilized every 2 weeks with  $100 \text{ mg L}^{-1}$  of Peter’s 20-20-20 N-P-K fertilizer.

#### Molecular characterization of hybrid plants

PCR was performed on seedlings from  $T_2$  hybrid families to identify transgenic and non-transgenic null segregant progeny from each cross. A leaf tip from each seedling was collected and total DNA was isolated using a CTAB buffer and two rounds of phenol/chloroform extraction followed by two ethanol washes. DNA was quantified using a BioTek plate analyzer and diluted to  $20 \text{ ng } \mu\text{L}^{-1}$ . PCR was performed using GoGreen Mastermix according to the manufacturers’ instructions using 80 ng of DNA. Two pairs of primers were used to verify transgene status (presence or absence of transgene). *Gus* linker primers (sequences in Fu et al. 2011) were used to confirm presence of the transgene, and *hyg* primers (sequences in Shen et al. 2012) were used to identify the hygromycin selection marker gene. PCR for each  $T_2$  hybrid family was performed with a positive control (the  $T_1$  COMT transgenic parent) and a negative control (the wild type field selection parent). Seedlings showing both amplification products were considered positive ( $T_2$  COMT transgenic), while those with no amplification products were considered negative (non-transgenic null segregant).

#### Morphology characterization

Morphological measurements were taken on seedlings at 40 and 80 DAS, and on plants of the first reproductive (R1) stage (developmental stages determined using the criteria described by Hardin et al. (2013). Early morphological traits (height at 40 and 80 DAS, number of tillers at 80 DAS) were measured on seedlings maintained in  $7.6 \times 7.6 \times 5.7 \text{ cm}$  containers. Height was measured from soil level to the tip of



**Fig. 1** Regressions between  $T_2$  hybrid family seed weight and **a** germination rate, **b** height at 40 days after sowing, **c** height at 80 days after sowing, and **d** tiller number at 80 days after sowing. Mean seed weight was determined for 11  $T_2$  hybrid

families; each data point represents an average of 15–18 biological replicates per family. Values in **bold** indicate significance at the 5 % level

the tallest leaf. At 80 DAS, plants were transferred to 12-L pots. Based on 80 day morphology measurements,  $T_2$  hybrids at the top 10 % or bottom 10 % of trait (plant height and tiller number 80 DAS) distribution tails were selected for further experiments. The middle 80 % of plants fell into the non-selected category.

Once tillers reached the R1 growth stage, morphological traits (tiller height, leaf width, and number of tillers) were measured on selected plants. After taking morphology measurements, dry biomass was determined for selected and non-selected plants. For non-selected plants, all biomass above

the soil level was harvested, dried in an oven at 43 °C for 96 h, and weighed. For selected plants, half of each plant was harvested for dry weight determination and the resulting weight was multiplied by two to estimate the total dry weight yield. The remaining half was maintained for possible future field experiments.

#### Cell wall chemistry and sugar release

Above-ground tillers at the R1 growth stage were harvested from selected and non-selected plants, dried in an oven at 43 °C for 96 h, and milled with

**Table 1** Correlations among morphological and cell wall variables measured on transgenic (shaded) and non-transgenic null segregant (non shaded) T<sub>2</sub> hybrid offspring

	40 day height	80 day tiller no.	80 day height	R1 height	R1 tiller no.	R1 leaf width	Dry weight	Lignin content	S/G ratio	Sugar release
40 day height		<b>0.59</b> <.0001	0.52 <.0001	<b>0.64</b> <.0001	<b>0.59</b> <.0001	0.20 0.17	0.47 <.0001	0.11 0.22	0.45 <.0001	-0.094 0.59
80 day tiller no.	0.52 <.0001		0.40 <.0001	<b>0.60</b> <.0001	<b>0.67</b> <.0001	0.25 0.093	0.47 <.0001	0.17 0.072	0.38 <.0001	-0.033 0.85
80 day height	<b>0.61</b> <.0001	0.36 <.0001		0.26 0.051	-0.047 0.74	-0.24 0.094	0.09 0.28	0.12 0.19	0.14 0.12	-0.21 0.22
R1 height	0.36 0.037	0.54 0.001	0.30 0.083		<b>0.75</b> <.0001	0.37 0.010	<b>0.74</b> <.0001	0.52 <.0001	<b>0.57</b> <.0001	-0.005 0.98
R1 tiller no.	0.51 0.002	<b>0.68</b> <.0001	0.22 0.21	0.58 <.0001		0.35 0.015	<b>0.91</b> <.0001	0.49 0.001	<b>0.57</b> <.0001	0.18 0.46
R1 leaf width	0.32 0.081	0.15 0.43	0.14 0.45	0.36 0.044	0.53 0.002		0.34 0.023	0.04 0.80	0.12 0.50	0.39 0.12
Dry weight	0.25 0.006	0.31 0.001	0.23 0.013	0.47 0.009	<b>0.72</b> <.0001	0.38 0.049		0.48 <.0001	<b>0.58</b> <.0001	0.062 0.73
Lignin content	0.063 0.54	0.15 0.15	0.21 0.035	-0.048 0.82	0.14 0.50	-0.043 0.85	0.29 0.004		0.42 <.0001	-0.43 0.008
S/G ratio	0.19 0.071	0.017 0.87	0.074 0.47	0.14 0.50	0.33 0.11	0.14 0.52	0.26 0.011	0.39 <.0001		-0.090 0.60
Sugar release	-0.028 0.88	-0.243 0.18	-0.038 0.84	-0.26 0.40	-0.23 0.46	0.039 0.90	-0.076 0.69	-0.26 0.15	-0.39 0.022	

Values are Pearson correlations with corresponding p values below. n = 163 for transgenic offspring, n = 198 for non-transgenic null segregant offspring. Values in bold represent the strongest ( $\geq 0.56$ ) correlations. Morphology measurements were taken at 40 and 80 days after sowing (40 day height, 80 day height and tiller number) and at the first reproductive stage (R1 height, tiller number, and leaf width). Lignin content, S/G (syringyl/guaiacyl) lignin monomer ratio, and sugar release were measured on whole tillers harvested at the R1 stage

a Wiley mill (Thomas Scientific, Model 4, Swedesboro, NJ, USA) through a 0.5 mm screen. Milled samples were used for lignin and sugar release analyses.


Pyrolysis mass beam mass spectrometry (py-MBMS) was used to measure total lignin content and the S/G lignin monomer ratio as previously described (Sykes et al. 2009). After the removal of soluble extractives and starch, the remaining cell wall residues (CWR) were pyrolyzed at 500 °C and resulting vapors were analyzed via MBMS. Lignin content was estimated as the sum of the relative intensities of all lignin precursor peaks. The S/G lignin monomer ratio was estimated as the sum of the intensity of syringyl peaks divided by the sum of the intensity of guaiacyl peaks.

The total sugars released by enzymatic hydrolysis were measured by high-throughput sugar release assays as previously described (Selig et al. 2010). Briefly, soluble extractives and starches were removed from samples (Decker et al. 2012) and resulting CWR were pretreated with condensing steam at 180 °C for

17.5 min, followed by an incubation with 70 mg protein/g biomass Ctec2 enzyme cocktail (Novozymes North America, Franklinton, NC, USA) at 50 °C for 70 h. Colorimetric assays were used to determine the amounts of glucose and xylose released by the samples (Megazyme Intl., Bray, Ireland). The total sugar release reported is the sum of the glucose and xylose release.

#### Data analysis

All data analysis was performed using SAS Statistical Analysis Software v.9.3 (SAS Institute, Cary, NC, USA). Analysis of variance (ANOVA) was used to partition variance in morphological and cell wall variables within and among families, parent types (COMT transgenic or field accession), family types (full-sibling or half-sibling, and progeny transgene status). Means were separated using Tukey's studentized range test. The REG and CORR procedures were used to perform single linear regressions and correlation analyses, respectively. Chi square goodness of fit



**Fig. 2** Effect of female parent type, T<sub>1</sub> COMT transgenic (+) or wild type (WT) field selection, on early morphological traits in T<sub>2</sub> hybrid families. **a** Height at 40 days after sowing, **b** height at 80 days after sowing, and **c** number of tillers at 80 days after sowing. Bars represent the mean of biological replicates for T<sub>2</sub> hybrid families generated with either T<sub>1</sub> COMT transgenic as the female parent (n = 14) or wild type field selection as the female parent (n = 16) ± standard error of the mean. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey's studentized range test. Bars with different letters are significantly different at the 5 % level (p < 0.05)

was performed using PROC FREQ with a hypothesized distribution of equal proportions of COMT (+) and COMT (−) offspring in each family.

## Results

Relationships among seed weight, germination rate, and early-growth morphological variables


Final germination was tallied at 30 DAS. Height at 40 and 80 DAS and number of tillers at 80 DAS was determined for T<sub>2</sub> hybrid families. Seed weight did not significantly influence germination rate, but positively influenced seedling height at 40 DAS ( $R^2 = 0.43$ ; Fig. 1b). All early morphological variables (height at 40 and 80 DAS and number of tillers at 80 DAS) were significantly positively correlated with one another, with Pearson correlation coefficients ranging from 0.36 to 0.61 (Table 1).

Effect of female parent type on morphology and cell wall traits

T<sub>1</sub> COMT × field accession hybrids exhibited morphological differences in early (40 and 80 DAS) and late (R1 stage) growth phases, which varied by female parent. T<sub>2</sub> hybrid families produced using the T<sub>1</sub> COMT transgenic line as the female parent were 39 % taller at 40 DAS and had 20 % more tillers at 80 DAS than those families produced using the non-transgenic field switchgrass as the female parent (Fig. 2). The T<sub>1</sub> COMT transgenic line female parent produced hybrids that were 10 % taller at the R1 stage, which produced 113 % more tillers and 63 % more biomass than hybrids coming from seed produced by the field clone (Fig. 3). Female parent type had no effect on lignin content, S/G ratios, or total sugar release in the T<sub>2</sub> hybrid families (Fig. 4).

Transgene segregation

T<sub>2</sub> hybrid families were genotyped for presence (+) or absence (−) of the COMT transgene. Overall, families had a mean of 47 % COMT (+) offspring. Distribution of COMT (−) and COMT (+) T<sub>2</sub> hybrid offspring was not statistically different from 50 % for most families (Table 2). Three families had more COMT (−) offspring than the expected 50 %, while two families had more COMT (+) offspring than expected. Analysis of variance showed that female parent type (T<sub>1</sub> COMT transgenic or wild type field selection) and the interaction of female parent type and family type




**Fig. 3** Effect of female parent type, T<sub>1</sub> COMT transgenic (+) or wild type (WT) field selection, on reproductive-stage morphology and dry weight yield in T<sub>2</sub> hybrid families. Measurements were taken once tillers reached the first reproductive stage (R1). **a** Tiller height, **b** leaf width, **c** number of tillers, and **d** dry weight yield. Bars represent the mean of biological replicates for T<sub>2</sub> hybrid families generated with either

T<sub>1</sub> COMT transgenic as the female parent (n = 14) or wild type field selection as the female parent (n = 16) ± standard error. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey's studentized range test. Bars with different letters are significantly different at the 5 % level (p < 0.05)

(full-sibling or half-sibling) were significant sources of variation in the percent of COMT (+) offspring in a family (Table 3). Specifically, the offspring of half-sibling families with a wild type female parent had only 20 % COMT (+) offspring, on average. All other 95 % confidence limits of percent COMT (+) offspring for female parent/family type combinations included 50 % (Table 4).

Effect of transgene status on morphology and cell wall traits

Transgenic seedlings were similar to the null segregants in height at 40 DAS, but were 7 % shorter by 80 DAS (Fig. 5). Once reaching the R1 developmental stage, there were no significant differences between the transgenic hybrids and null segregants in height,



**Fig. 4** Effect of female parent type, T<sub>1</sub> COMT transgenic (+) or wild type (WT) field selection, on cell wall traits in T<sub>2</sub> hybrid families. Whole tillers of the first reproductive stage (R1) were used for cell wall analyses. **a** MBMS lignin content, **b** S/G lignin monomer ratio, and **c** total sugars (glucose and xylose combined) released by enzymatic hydrolysis. Bars represent the mean of biological replicates for T<sub>2</sub> hybrid families

leaf width, or tiller number. However, transgenic hybrids had 43 % less dry weight compared with null segregants (Fig. 6).

As expected, there were clear differences in cell wall chemistry and recalcitrance between the T<sub>2</sub> transgenic hybrids and the non-transgenic null segregants. R1-stage tillers harvested from COMT transgenic hybrids had 7 % less lignin than null segregants, as well as a 12 % reduction in the S/G ratio. Changes in lignin content and composition were accompanied by a 12 % increase in total sugar release (Fig. 7).

#### Genetic control of morphology and cell wall components

Genetics had a clear influence on T<sub>2</sub> hybrid switchgrass morphology. There were significant differences among families for early morphological variables (height at 40 and 80 DAS and tiller number at 80 DAS), R1 stage morphological variables (height, tiller number, and leaf width) and dry weight yield (Table 5). There was a 6-fold difference in the shortest and tallest families at 40 DAS. At 80 DAS, the difference between the shortest and tallest families was only 1.5-fold (Fig. 8). Families were ranked based on height at 40 and 80 days; spearman rank correlation indicated that 68 % of the variation in 80 day height could be explained by 40 day height ( $r_s = 0.68$ ,

generated with either T<sub>1</sub> COMT transgenic as the female parent ( $n = 14$ ) or wild type field selection as the female parent ( $n = 16$ )  $\pm$  standard error. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey's studentized range test. Bars with different letters are significantly different at the 5 % level ( $p < 0.05$ ). CWR, cell wall residues; (+), transgenic; WT, wild type

$p < 0.001$ ). There were also some differences among families for cell wall variables (lignin content, S/G ratio, and total sugar release), although, transgene status was much more influential than family for these traits. Conversely, family was more influential than transgene status for growth-related variables (early and R1 stage morphology, dry weight yield; Table 5).

Distribution of transgenic hybrid offspring among the T<sub>2</sub> switchgrass families genotyped ranged from 0 to 100 % (Table 2). In order to differentiate between family (genetic) effects and effects resulting from uneven transgene segregation, we regressed family means for each morphological and cell-wall variable on percent transgenic offspring for each family. There was no significant influence of percent transgenic offspring on any of the morphological variables, cell-wall variables, or family rankings (Table 6). We conclude that family effects on morphological and cell-wall traits can be interpreted as the result of genetic differences between families rather than differences related to uneven transgene segregation between families.

#### Correlations among morphology, dry weight yield, and cell wall variables

Many morphological and cell wall variables were significantly correlated (Table 1). Most of the early



**Table 2** Transgene segregation for T<sub>2</sub> hybrid switchgrass families

Family Number	No. seedlings genotyped	Female parent type	Family type	Percent COMT(+) offspring	$\chi^2$ value
4	14	C	FS	50	0
6	18	C	FS	39	0.89
12	15	C	FS	80	5.4**
13	17	C	FS	53	0.05
16	15	C	FS	33	1.6
18	14	C	FS	36	1.1
20	8	C	FS	50	0
23	14	C	FS	36	1.1
24	14	C	FS	64	1.1
1	9	C	HS	11	5.4**
8	12	C	HS	67	1.3
27	13	C	HS	62	0.69
29	15	C	HS	53	0.067
30	14	C	HS	100	N/A
2	9	W	FS	78	2.7
3	16	W	FS	25	4.0**
7	12	W	FS	42	0.33
9	15	W	FS	27	3.2
10	16	W	FS	63	1
11	17	W	FS	42	0.53
14	12	W	FS	42	0.33
15	12	W	FS	50	0
17	14	W	FS	50	0
22	15	W	HS	0	N/A
26	9	W	HS	33	1
28	12	W	HS	25	3.0

Families are ordered by female parent type and family type

Female parent type: *C* T<sub>1</sub> COMT transgenic, *W* wild type field selection; Family type: *FS* full-sibling, *HS* half-sibling; *Chi square value* Chi square goodness of fit test statistic based on a hypothetical distribution of equal percentages of COMT transgenic (+) and non-transgenic T<sub>2</sub> hybrid offspring; *N/A* Chi square testing was not possible due to an observation of zero (0) in a category

\*\* Chi square goodness of fit test statistics significant at the  $\alpha = 0.05$  level

**Table 3** Analysis of variance for influence of female parent type and family type on percent transgenic offspring in T<sub>2</sub> hybrid switchgrass families

Source	df	Type III SS	Mean square	F Value	Pr > F	Model R <sup>2</sup>
Female parent type <sup>a</sup>	1	2302.7	2302.7	5.8	0.03	0.26
Family type <sup>b</sup>	1	411.0	411.0	1.0	0.32	
Female parent $\times$ family type	1	1794.5	1794.5	4.5	0.05	
Error	22	8776.1	389.9			

<sup>a</sup> Female parent type (T<sub>1</sub> COMT transgenic or wild type field selection)

<sup>b</sup> Family type (full-sibling or half-sibling)

**Table 4** Mean percent of transgenic T<sub>2</sub> hybrid offspring with 95 % confidence limits for female parent type (T<sub>1</sub> COMT transgenic or wild type field selection) by family type (full-sibling or half-sibling) combinations

Female parent	Family type	n	Percent transgenic offspring	95 % confidence limits
T <sub>1</sub> COMT transgenic	FS	9	49.0 ± 6.7	35.2–62.8
T <sub>1</sub> COMT transgenic	HS	5	58.6 ± 8.9	40.1–77.1
Wild type	FS	9	46.5 ± 6.7	32.7–60.4
Wild type	HS	3	19.3 ± 11.5**	0–43

Family type FS full-sibling, HS half-sibling

(40 and 80 DAS) and R1 stage morphological variables were positively correlated with one another. In general, COMT (–) offspring showed more and larger significant trait correlations than COMT (+) offspring. Correlations between height at 40 DAS and height at R1 stage, and height at 40 DAS and dry weight, were twice as strong for COMT (–) offspring than for COMT (+) offspring. Similarly, correlations of dry weight with lignin content and S/G ratio were 0.48 and 0.58, respectively, for COMT (–) offspring, while the same correlations were 0.29 and 0.26 for COMT (+) offspring. There were no significant correlations between R1 morphological traits (height and tiller number) and cell wall traits (lignin and S/G ratio) for COMT (+) offspring, while the same correlations in COMT (–) offspring were significant. Lignin content and sugar release showed a predictable significant negative correlation in COMT (–) offspring, while there was no significant relationship between lignin content and sugar release for COMT (+) offspring.

## Discussion

Combining genetic engineering and breeding to incorporate reduced recalcitrance and other key biofuel traits into selected locally-adapted varieties could be a promising strategy for developing biofuel feedstocks that are high-yielding, well-adapted to their environment, and more amenable to bioconversion. To test the feasibility of this approach in a transgenic switchgrass line, we crossed low-lignin COMT-down-regulated switchgrass lines with high-yielding varieties selected from local Tennessee fields and characterized growth and recalcitrance among the resulting T<sub>2</sub> hybrid progeny. These results should provide baseline information to facilitate future


biofuel-related breeding efforts with transgenic switchgrass.

### Relationships among seed weight, germination rate, and early morphology

Seed weight is an agronomically important trait as it has been shown to be positively associated with higher germination rate, improved emergence, early seedling vigor, and faster growth rates (Das et al. 2004). We examined relationships among seed weight, germination rate, and early morphological variables in T<sub>2</sub> hybrid families. Although there was no significant effect of seed weight on germination rate in our study, seed weight positively influenced seedling height at 40 DAS (Fig. 1b), suggesting that seed weight does influence the initial growth rate (from day 0 to day 40) of young switchgrass seedlings. However, it should be noted that these experiments were performed under optimized greenhouse conditions; therefore, the results might not fully reflect the influence that seed weight would have on early growth traits if grown in a natural field setting where stresses are present.

### Effects of female parent type

The use of one particular genetic background over another as the female parent can dramatically influence growth and yield-related traits in hybrid offspring (Bhandari et al. 2014). In our study, there were no morphological or cell wall differences between T<sub>2</sub> hybrid switchgrass families with an ‘Alamo’ female parent and those with a ‘Kanlow’ female parent. However, female parent type (T<sub>1</sub> COMT transgenic or field accession) strongly influenced growth but had no effect on cell wall traits among the T<sub>2</sub> hybrids. In general, the T<sub>1</sub> COMT transgenic line used as the female parent had positive effects on growth



**Fig. 5** Effect of transgene status, absence (–) or presence (+) of transgene, on early morphological traits in T<sub>2</sub> hybrid families. **a** Height at 40 days after sowing, **b** height at 80 days after sowing, and **c** number of tillers at 80 days after sowing. Bars represent the mean of biological replicates for non-transgenic null segregant (n = 172) and T<sub>2</sub> COMT transgenic hybrid families (n = 140) ± standard error. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey’s studentized range test. Bars with different letters are significantly different at the 5 % level (p < 0.05)

morphology in the T<sub>2</sub> hybrid families, including improved height at early and late stages of growth, significantly more tillering, and a 63 % increase in dry weight yield over those hybrids produced using the wild type field selection as the female parent


(Figs. 2, 3). It might not necessarily be the use of a transgenic plant as a female that leads to improved growth, but that the transgenics used in this study were in a more favorable genetic background compared to that of the field selections. However, it is important to emphasize that these studies were performed under controlled greenhouse conditions. It is possible that the field-selected wild types could become the favored female parent if the T<sub>2</sub> hybrid families were grown under natural field conditions.

#### Transgene effects

Greenhouse-grown T<sub>1</sub>-generation COMT-downregulated switchgrass had reduced lignin, lower S/G lignin monomer ratios, and improved ethanol yield compared with non-transgenic null segregant controls (Fu et al. 2011). In our study, the altered lignin phenotype of the T<sub>1</sub> COMT parent was successfully transmitted to the T<sub>2</sub> transgenic hybrid families as demonstrated by the reduction in lignin content, lower S/G ratio, and higher sugar release relative to the null segregant progeny (Fig. 7). Despite morphological differences among the T<sub>2</sub> hybrid families, the strong effect of transgene status relative to family for lignin content, S/G ratio, and sugar release (Table 5) reiterates the strong genetic control of cell wall chemistry provided by the down-regulated *COMT* gene. Conversely, the strong effect of family relative to that of transgene status for morphological variables indicates that high-performing families can be distinguished before genotyping for presence or absence of the transgene. Genotyping only the top-performing families and individuals saves time and cost in plant breeding programs where narrowing down the number of individuals to maintain and evaluate at early stages is paramount to efficient genetic gain.

#### Relationships among morphology, dry weight yield, and cell wall traits

Relationships among early- and late- stage morphological traits, dry weight yield, and cell wall traits were analyzed for transgenic and non-transgenic T<sub>2</sub> hybrid families. In general, these data indicated that the non-transgenic null segregant offspring show stronger and more predictable patterns of plant growth, and more predictable relationships between morphology and cell-wall chemistry, than do transgenic offspring. For




**Fig. 6** Effect of transgene status, absence (–) or presence (+) of transgene, on reproductive-stage morphology and dry weight yield in T<sub>2</sub> hybrid families. Measurements were taken once tillers reached the first reproductive stage (R1). **a** Tiller height, **b** leaf width, **c** number of tillers, and **d** dry weight yield. Bars represent the mean of biological replicates for non-transgenic

null segregant ( $n = 46$ ) and T<sub>2</sub> COMT transgenic hybrid families ( $n = 42$ )  $\pm$  standard error. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey's studentized range test. Bars with different letters are significantly different at the 5 % level ( $p < 0.05$ )

both transgenic and non-transgenic offspring, there were significant positive correlations among most early (40 and 80 DAS) and mature (R1 stage) morphological traits, which would suggest that initial selections of hybrid switchgrass families with favorable growth performance could be performed at early

stages of development. Dry weight was significantly positively correlated with most morphological variables (Pearson correlation coefficients ranging from 0.23 to 0.91), but was most strongly correlated with R1 tiller number ( $r = 0.72$  for transgenic offspring,  $r = 0.91$  for non-transgenic offspring) followed by



**Fig. 7** Effect of transgene status, absence (–) or presence (+) of transgene, on cell wall traits in T<sub>2</sub> hybrid families. Whole tillers of the first reproductive stage (R1) were used for cell wall analyses. **a** MBMS lignin content, **b** S/G lignin monomer ratio, and **c** total sugar (glucose and xylose) released by enzymatic hydrolysis. Bars represent the mean of biological replicates for

non-transgenic null segregant (n = 104) and T<sub>2</sub>-COMT transgenic hybrid families (n = 116) ± standard error. Means were analyzed with a one-way ANOVA and letter groupings were obtained using Tukey’s studentized range test. Bars with different letters are significantly different at the 5 % level (p < 0.05). CWR cell wall residues

**Table 5** Influence of family and transgene status (presence or absence of transgene) on morphological and cell wall variables in T<sub>2</sub> hybrid switchgrass families

Source	df	40 Day height	80 Day height	80 Day tiller no.	R1 height	R1 leaf width	R1 tiller no.	Dry weight	Lignin content	S/G ratio	Sugar release
Family	27	10.26**	3.31**	7.34**	6.1**	2.37**	5.09**	8.53**	4.35**	5.95**	2.54**
Transgene status	1	0.15	0.93	0	1.68	0.54	0.4	7.98**	38.12**	74.31**	31.21**
Model R <sup>2</sup>		0.60	0.33	0.51	0.62	0.39	0.59	0.49	0.46	0.56	0.71

Means were separated using Tukey’s studentized range test. Asterisks represent F values significant at the 5 % level. Measurements were taken at 40 and 80 days after sowing, and at the first reproductive stage (R1)


R1 height (r = 0.47 for transgenic offspring, r = 0.74 for non-transgenic offspring). This is in agreement with prior studies that have found switchgrass biomass yield to be most strongly correlated with tillering and/or height compared to other morphological variables (Bhandari et al. 2010; Das et al. 2004). Taken together, these results suggest that tiller height and tiller number could be used as a selection index for identifying potentially high-yielding hybrid switchgrass families for further analyses.

Interestingly, there were no significant correlation between R1 morphological traits (height, tiller number) and cell wall traits (lignin, S/G ratio) for transgenic offspring. However, the same correlations in the non-transgenic null segregant offspring were

highly significant (Table 1). This suggests that in non-transgenic switchgrass, favorable plant morphology and high lignin content may be inherited together to some degree. In a breeding program, this would be an important consideration when the goal is to select for families that exhibit reduced recalcitrance (low lignin content) but still display desirable growth characteristics.

In the original greenhouse study, the desired traits were passed on to the T<sub>1</sub>-generation transgenic offspring with no apparent growth effects when compared to the corresponding non-transgenic null segregants (Fu et al. 2011), and these results were reproducible under field conditions in fully-established plants (Baxter et al. 2014). In the current study,

**Fig. 8** Least squares means for 40 day height, 80 day height, and 80 day tiller number for  $T_2$  hybrid switchgrass families. Families are ordered by 40 day height. Error bars represent standard deviation ( $n = 8\text{--}18$  seedlings/family). Measurements were taken at 40 and 80 days after sowing



**Table 6** Summary of single linear regressions of family ranks and mean morphological and cell wall variables on percent of  $T_2$  COMT transgenic offspring in each family

Independent variable	<i>df</i>	<i>MS</i>	<i>F</i>	<i>Pr &gt; F</i>	Model <i>R</i> <sup>2</sup>
40 DAS rank	1	31.1	0.42	0.52	0.02
40 DAS height	1	47.8	1.10	0.30	0.04
80 DAS rank	1	39.1	0.56	0.46	0.02
80 DAS height	1	51.4	2.57	0.12	0.09
80 DAS tiller number	1	1.2	3.61	0.07	0.12
R1 height	1	716.7	1.17	0.29	0.06
R1 tiller number	1	36.4	0.54	0.47	0.00
R1 leaf width	1	0.0	0.02	0.90	0.00
Dry weight	1	26.8	0.04	0.84	0.01
S/G ratio	1	0.00	1.94	0.18	0.07
Lignin content	1	0.11	0.11	0.74	0.00

Morphology measurements were taken at 40 and 80 days after sowing (DAS), and at the first reproductive stage (R1). Lignin content and S/G ratio were determined for R1-stage whole tillers

COMT-downregulation in the  $T_2$  generation does appear to affect growth in the transgenic hybrids when compared with the non-transgenic null segregants, as reflected by the decreased dry weight yield (43 %). In our case, the goal is to increase biomass yield relative to the transgenic parent while maintaining the desired cell wall traits. Thus, the ideal  $T_2$  hybrid family would exhibit low recalcitrance while producing biomass amounts at least equal to their  $T_1$  transgenic parent, not necessarily the corresponding  $T_2$  null segregant. Multi-year field experiments will provide an opportunity to compare morphology, biomass yield, and

recalcitrance between  $T_2$  hybrid offspring and their  $T_1$  transgenic parent in a realistic environment, and to examine potential tradeoffs between agronomic properties and cell wall recalcitrance.

**Acknowledgments** We thank Fred Allen, Hem Bhandari, and Ken Goddard for providing the Tennessee field accessions of switchgrass field selections, and Justin Leduc and Erika Barton for their assistance with collecting and preparing samples for cell wall characterization analyses. This work was supported by the Agriculture and Food Research Initiative (United States Department of Agriculture) and Southeastern Partnership for Integrated Biomass and Supply Systems (The IBSS Partnership), and enabled by the Bioenergy Science Center. The Bioenergy

Science Center is a U.S. Department of Energy Bioenergy Research Center supported by the Office of Biological and Environmental Research in the DOE Office of Science.

### Compliance with ethical standards

**Conflict of interest** The authors declare that they have no competing interests.

### References

- Baxter HL, Mazarei M, Labbé N, Kline LM, Cheng Q, Windham M, Stewart CN Jr (2014) Two-year field analysis of reduced recalcitrance transgenic switchgrass. *Plant Biotechnol J* 12:914–924
- Baxter HL, Poovaiah CR, Yee KL, Mazarei M, Rodriguez M Jr, Thompson OA, Shen H, Turner G, Decker SR, Sykes RW, Chen F, Davis MF, Mielenz JR, Davison BH, Dixon RA, Stewart CN Jr (2015) Field evaluation of transgenic switchgrass plants overexpressing PvMYB4 for reduced biomass recalcitrance. *Bioenergy Res* 8:910–921
- Bhandari HS, Saha MC, Mascia PN, Fasoula VA, Bouton JH (2010) Variation among half-sib families and heritability for biomass yield and other traits in lowland switchgrass. *Crop Sci* 50:2355–2363
- Bhandari HS, Webb SL, Bouton JH, Saha MC (2014) Reciprocal effects for biomass yield in lowland switchgrass. *Crop Sci* 54:955–962
- Boerjan W, Ralph J, Baucher M (2003) Lignin biosynthesis. *Annu Rev Plant Biol* 54:519–546
- Chen F, Dixon RA (2007) Lignin modification improves fermentable sugar yields for biofuel production. *Nat Biotechnol* 25:759–761
- Das MK, Fuentes RG, Taliaferro CM (2004) Genetic variability and trait relationships in switchgrass. *Crop Sci* 44:443–448
- Decker SR, Carlile M, Selig MJ, Doepcke C, Davis M, Sykes R et al (2012) Reducing the effect of variable starch levels in biomass recalcitrance screening. In: Himmel ME (ed) *Biomass conversion*. Humana Press, New York, pp 181–195
- Fu C, Mielenz JR, Xiao X, Ge Y, Hamilton CY, Rodriguez M Jr, Chen F, Foston M, Ragauskas A, Bouton J, Dixon RA, Wang ZY (2011) Genetic manipulation of lignin reduces recalcitrance and improves ethanol production from switchgrass. *Proc Natl Acad Sci USA* 108:3803–3808
- Fu C, Sunkar R, Zhou C, Shen H, Zhang JY, Matts J, Wang ZY (2012) Overexpression of miR156 in switchgrass (*Panicum virgatum* L.) results in various morphological alterations and leads to improved biomass production. *Plant Biotechnol J* 10:443–452
- Ge Y, Fu C, Bhandari H, Bouton J, Brummer EC, Wang ZY (2011) Pollen viability and longevity of *switchgrass* (L.). *Crop Sci* 51:2698–2705
- Hardin CF, Fu C, Hisano H, Xiao X, Shen H, Stewart CN, Wang ZY (2013) Standardization of switchgrass sample collection for cell wall and biomass trait analysis. *Bioenergy Res* 6:755–762
- McLaughlin SB, Walsh ME (1998) Evaluating environmental consequences of producing herbaceous crops for bioenergy. *Biomass Bioenergy* 14:317–324
- Parrish DJ, Fike JH (2005) The biology and agronomy of switchgrass for biofuels. *BPTS* 24:423–459
- Poovaiah CR, Nageswara-Rao M, Soneji JR, Baxter HL, Stewart CN (2014) Altered lignin biosynthesis using biotechnology to improve lignocellulosic biofuel feedstocks. *Plant Biotechnol J* 12:1163–1173
- Poovaiah CR, Mazarei M, Decker SR, Turner GB, Sykes RW, Davis MF, Stewart CN Jr (2015) Transgenic switchgrass (*Panicum virgatum* L.) biomass is increased by overexpression of switchgrass sucrose synthase (PvSUS1). *Biotechnol J* 10:552–563
- Sarath G, Akin DE, Mitchell RB, Vogel KP (2008) Cell-wall composition and accessibility to hydrolytic enzymes is differentially altered in divergently bred switchgrass (*Panicum virgatum* L.) genotypes. *Appl Biochem Biotechnol* 150:1–14
- Schmer MR, Vogel KP, Mitchell RB, Perrin RK (2008) Net energy of cellulosic ethanol from switchgrass. *Proc Natl Acad Sci USA* 105:464–469
- Selig MJ, Tucker MP, Sykes RW, Reichel KL, Brunecky R, Himmel ME, Davis MF, Decker SR (2010) Biomass recalcitrance screening by integrated high throughput hydrothermal pretreatment and enzymatic saccharification. *Ind Biotechnol* 6:104–111
- Shen H, He X, Poovaiah CR, Wuddineh WA, Ma J, Mann DG, Wang H, Jackson L, Tang Y, Stewart CN Jr, Chen F, Dixon RA (2012) Functional characterization of the switchgrass (*Panicum virgatum*) R2R3-MYB transcription factor PvMYB4 for improvement of lignocellulosic feedstocks. *New Phytol* 193:121–136
- Sykes R, Yung M, Novaes E, Kirst M, Peter G, Davis M (2009) High-throughput screening of plant cell-wall composition using pyrolysis molecular beam mass spectroscopy. In: Mielenz JR (ed) *Biofuels: methods and protocols*, methods in molecular biology. Humana Press, New York, pp 169–183
- Vogel KP, Mitchell RB, Sarath G, Jung HG, Dien BS, Casler MD (2013) Switchgrass biomass composition altered by six generations of divergent breeding for digestibility. *Crop Sci* 53:853–862
- Wuddineh WA, Mazarei M, Zhang J, Poovaiah CR, Mann DG, Ziebell A, Sykes RW, Davis MF, Udvardi MK, Stewart CN Jr (2015a) Identification and overexpression of gibberellin 2-oxidase (GA2ox) in switchgrass (*Panicum virgatum* L.) for improved plant architecture and reduced biomass recalcitrance. *Plant Biotechnol J* 13:636–647
- Wuddineh WA, Mazarei M, Turner GB, Sykes RW, Decker ST, Davis MF, Stewart CN Jr (2015b) Identification and molecular characterization of the switchgrass AP2/ERF transcription factor superfamily, and overexpression of PvERF001 for improvement of biomass characteristics for biofuel. *Front Bioeng Biotechnol* 3:101. doi:10.3389/fbioe.2015.00101